



Requirement of LaeA for secondary metabolism and sclerotial production in *Aspergillus flavus*

Shubha P. Kale^a, Lane Milde^b, Marisa K. Trapp^b, Jens C. Frisvad^c, Nancy P. Keller^{b,d,*}, Jin Woo Bok^b

^a Department of Biology, Xavier University of Louisiana, New Orleans, LA 70125, USA

^b Department of Plant Pathology, University of Wisconsin-Madison, 882 Russell Labs, 1630 Linden Drive, Madison, WI 53706, USA

^c Center for Microbial Biotechnology DTU Biosys, Technical University of Denmark, Lyngby, Denmark

^d Department of Medical Microbiology and Immunology, University of Wisconsin-Madison, Madison, WI 53706, USA

ARTICLE INFO

Article history:

Received 19 May 2008

Accepted 27 June 2008

Available online 11 July 2008

Keywords:

Aflatoxin

LaeA

AflR

VeA

Sclerotia

Mycotoxin

Secondary metabolism

ABSTRACT

The nuclear regulator LaeA has been shown to govern production of multiple secondary metabolites in *Aspergillus nidulans* and *Aspergillus fumigatus*. Herein we examine the role of this protein in *Aspergillus flavus*. Similarly as in other Aspergilli, LaeA had a major effect on *A. flavus* secondary metabolism where $\Delta laeA$ and over-expression laeA (OE::laeA) strains yielded opposite phenotypes resulting in decreased (increased) secondary metabolite production. The two mutant strains also exhibited striking morphological phenotypes in the loss (increase) of sclerotial production in comparison to wildtype. Growth on seed was marked by decreased (increased) conidial and aflatoxin production of the respective mutants; this was accompanied by decreased lipase activity in $\Delta laeA$, an enzymatic process correlated with seed maceration. Transcriptional examination of the mutants showed LaeA negatively regulates expression of its recently identified nuclear partner VeA, another global regulator of *A. flavus* secondary metabolites and sclerotia.

© 2008 Elsevier Inc. All rights reserved.

1. Introduction

The genus *Aspergillus* represents a large, worldwide family of fungi with estimates nearing 200 species (Samson, 1992). Whereas most of the members of this genus are saprobes capable of thriving on plant, animal and manmade wastes, a few are potent pathogens of plants and animals. The two most frequently isolated pathogens from these diverse hosts are *Aspergillus flavus* and *Aspergillus fumigatus*, respectively. *A. flavus* is notorious for the production of the carcinogenic and mutagenic secondary metabolites commonly known as aflatoxins during growth on seed (Payne and Brown, 1998). In the US alone, agricultural economic losses due to aflatoxin contamination of food and feed are estimated to be \$270 million annually (Richard and Payne, 2003). However, the effects are more keenly felt in developing countries where, in addition to contributing to enhanced cancer rates (Bressac et al., 1991), high aflatoxin contamination of human food also contributes to fatal toxicoses (Probst et al., 2007). This species is also associated with invasive aspergillosis (IA) in immunocompromised individuals

* Corresponding author. Address: Department of Plant Pathology, University of Wisconsin-Madison, 882 Russell Labs, 1630 Linden Drive, Madison, WI 53706, USA. Fax: +1 608 263 2626.

E-mail addresses: skale@xula.edu (S.P. Kale), Irmilde@wisc.edu (L. Milde), trapp@plantpath.wisc.edu (M.K. Trapp), jcf@biocentrum.dtu.dk (J.C. Frisvad), npk@plantpath.wisc.edu (N.P. Keller), jbok@plantpath.wisc.edu (J.W. Bok).

(Hedayati et al., 2007). By far the majority of IA is attributed to *A. fumigatus* infection with instances of 50–90% mortality (Latge, 1999).

In common with all Aspergilli sequenced to date, both opportunistic pathogens have been found to contain a plethora of clustered genes devoted to the production of secondary metabolites or natural products. Many of these clusters were found to be regulated by a single nuclear protein called LaeA in *A. fumigatus* and the genetic model *A. nidulans* (Bok and Keller, 2004; Perrin et al., 2007). Deletion of *A. nidulans laeA* resulted in the reduction of secondary metabolite gene expression concomitant with loss/decrease in the production of the encoded metabolites, whereas overexpression of *laeA* yielded opposite results (Bok and Keller, 2004; Bok et al., 2006b). Deletion of the gene in two *A. fumigatus* strains resulted in decreased virulence in the mouse pulmonary model, associated with reduced killing of neutrophil cells (Bok et al., 2005; Sugui et al., 2007a) thus strongly supporting a role for LaeA mediated toxin production in IA development by *A. fumigatus*.

Here we examined the role of LaeA in the seed colonizing fungus *A. flavus*. Predictably, the $\Delta laeA$ mutant was crippled in secondary metabolite production whereas the overexpression strain overproduced several metabolites including the carcinogen aflatoxin. Furthermore, $\Delta laeA$ showed decreased lipase activity and was less able to colonize peanut and maize seed. Unexpectedly, however, $\Delta laeA$ was unable to produce sclerotia, overwintering structures important in fungal survival. Sclerotial and secondary metabolism

loss have also been reported in *ΔveA* mutants of *A. flavus* (Duran et al., 2007). VeA and LaeA have recently been identified as members of a nuclear complex in *A. nidulans* (Bayram et al., 2008) and regulation of *veA* expression by LaeA as shown here may represent an internal feedback mechanism to maintain a nuclear complex linking morphology and secondary metabolism in the aspergilli.

2. Materials and methods

2.1. Strains and growth conditions

All strains were maintained as glycerol stocks and grown on potato dextrose agar (PDA) or glucose minimal medium (GMM, Shimizu and Keller, 2001) for spore production at 29 °C. Media were supplemented with 10 mM uridine and 10 mM uracil as needed. Strains used in this study included *A. flavus* NRRL 3357 (prototroph), NRRL 3357.5 (*pyrG* auxotroph) and the transformants described below.

2.2. Vector construction

The *laeA* replacement vector was constructed as follows. A 1.4 kb fragment upstream of the *laeA* coding region was amplified with forward primer 5' TGTGTCGACACTGCCAGACATCTATA adding a Sall site as indicated by underline and reverse primer 5' GTAGTACGAGTCGTGTGGTGGTCCGGCCGCGC adding a NotI site as indicated. This fragment was digested with NsiI (which was internal to the Sall site) and NotI and ligated into the NsiI and NotI sites of pLMH26 (Maggio-Hall and Keller, unpublished data) upstream of the *A. fumigatus pyrG*. Next, a 1.5 kb fragment downstream of the *laeA* coding region was amplified with forward primer 5' CAGCCGCGGACGATGCACTGAGCTGCCT adding a SacII site and reverse primer 5' CCTCGCCAGCAACGGCCGAGACC approximately 20 bp downstream of a SacII site. This fragment was digested with SacII and ligated into pLMH26 linearized with SacII to yield pLRM5. The correct orientation of the downstream fragment was confirmed by PstI digestion and the correct candidates were then confirmed by digests with NotI and NsiI. The schematic of this replacement vector is depicted in Fig. 1A. The *laeA* complementation vector was constructed by amplifying a 4.3 kb fragment containing the 2 kb *laeA* coding region and 2.3 kb upstream with forward primer 5' GGGATCCTCCACAAAGCCTTTCGTAAAA and reverse primer 5' TATCTAGAAGCACAGGCATGCGGCCGCA. This fragment was ligated into pCR-Blunt-II Topo (Invitrogen) to create intermediate vector pLRM9. A 3 kb HindIII fragment containing the *niaD* gene was cut from pGAPN-2 (Liang et al., 1997) and ligated into the HindIII site of pLRM9 to create the *laeA* complementation vector pLRM11.

2.3. Fungal transformation

Aspergillus protoplasts were produced and transformed using the modified polyethylene glycol method (Bok and Keller, 2004). To generate *laeA::pyrG* (i.e. *ΔlaeA*) strains, *A. flavus* NRRL 3357.5 protoplasts were transformed with 2 μg of a 5 kb pLRM5 PCR product generated using the forward primer 5' CCTTGATGATGATGTA TGATGAGC and the reverse primer 5' TCTTGGTCATTGGGTGGCGG. Transformants were screened for uridine and uracil prototrophy followed by Southern analysis. The resulting *laeA* deletion strains used for further experiments were named as TJW71.1, TJW71.3, TJW71.4 and TJW71.7, respectively.

To generate nitrate auxotrophs of the *laeA* deletion strain, 10⁶ conidia from TJW71.1 were spread onto GMM containing 750 mM chlorate and 1.6 mM ammonium chloride. Colonies that arose were transferred to fresh plates and single spores were purified. The resulting colonies were screened for mutation of *niaD* as

described in Cove, where growth on different nitrogen sources allowed for classification of mutations in *niaD*, *niaA* and other genes required for nitrate utilization (Cove, 1966). One *niaD* auxotroph strain, MLRM8.1, was transformed with 10 μg of pLRM11 to yield the complemented strains, TJW79 series.

2.4. Physiology experiments

Conidial production, sclerotial formation and relative colony diameter were recorded for wildtype, *ΔlaeA* and complemented *laeA* strains. Diameter growth was measured from point inoculation of 5 μl of a 10⁶ spores/ml suspension of *A. flavus* conidia on GMM media. For quantitative analysis of conidial production, YES medium (2% yeast extract, 6% sucrose, pH 5.8) and GMM were overlaid with 5 ml of a 10⁶ spores/ml suspension of *A. flavus* conidia in molten agar. Cultures were grown for 5 days at 29 °C in light. Three 1.5-cm diameter cores were harvested from the center of each plate and homogenized in 3 ml of distilled water. 2 μl were removed, diluted 1:500, and conidia were counted using a hemocytometer. Sclerotial production was observed on the sclerotial inducing medium GMM + 2% sorbitol. GMM + 2% sorbitol was overlaid with 3 ml of a 10⁶ spores/ml suspension of *A. flavus* conidia in molten agar. Cultures were grown for 5 days at 29 °C in dark conditions. Plates were then sprayed with 70% ethanol to kill and wash away conidia to aid in enumeration of sclerotia.

3. Results

3.1. Identification and disruption of *A. flavus laeA*

The predicted sequence for the *A. flavus laeA* ortholog was obtained from GenBank (AY883016) and by conducting a BLAST search (<http://www.ncbi.nlm.nih.gov/blast/>) with the *A. nidulans* sequence against the *A. flavus* genome scaffolds (<http://www.aspergillusflavus.org/genomics/>) and then selecting the matching genomic region as well as 4 kb of up and downstream of the predicted *laeA* ORF using the gene prediction program FGENESH (<http://www.bio.net/bionet/mm/bio-www/1999-January/000775.html>). The resulting two exon gene was similar in structure to *A. nidulans* and *A. fumigatus laeA*. The *A. flavus laeA* gene is 100% identical to a putative *A. oryzae laeA* and its 5' exon aligns well with the sequences from *A. nidulans* and *A. fumigatus* with the most variation in the 5' region (data not shown). *A. flavus* LaeA protein shows 75.6% identity to *A. nidulans* LaeA and contains a conserved s-adenosyl methionine binding (SAM) motif found to be critical for LaeA function (Bok et al., 2006c). Inactivation of the *laeA* locus was obtained by replacement of *laeA* by *pyrG* (Supplementary Fig. 1A). Four out of 20 transformants contained the 4 and 1 kb fragments of a *SpeI* digest expected for a *laeA* disruption event and were called TJW71.1, TJW71.3, TJW71.4 and TJW71.7, respectively (Supplementary Fig. 1B). Physiological and pathogenicity experiments as below revealed all *ΔlaeA* strains shared the same phenotype and TJW71.1 was selected for the majority of experimentation.

A *niaD*-auxotroph of TJW71.1 was created to generate strain MLRM8.1 which was then complemented to *laeA* and *niaD* prototrophy with pLRM11. Four transformants were confirmed to contain wildtype *laeA* (Supplementary Fig. 1C and data not shown). Examination of a representative transformant strain, TJW79.13 containing at least two copies of *laeA* (Supplementary Fig. 1C), was selected for further experiments to examine effects of over expression *laeA*. This strain is referred to as OE::*laeA* in this study.




Fig. 1. Aflatoxin production and gene expression in $\Delta laeA$ and OE::laeA mutants. (A) Extracts of the wildtype *A. flavus* NRRL 3357, the $\Delta laeA$ mutant (TJW71.1) and four OE::laeA strains were grown on YES media for 5 days at 29 °C and extracts were separated on a thin layer chromatography plate. Each strain was extracted twice. Aflatoxin was visualized using long-wave (366 nm) UV light. Aflatoxin B1 standard is spotted on each side of the plate. (B) mRNA expression of *laeA*, *veA* and *aflR* in *A. flavus* NRRL 3357, $\Delta laeA$ (TJW71.1) and OE::laeA (TJW79.13). All strains were cultured in liquid YEP medium and incubated at 250 rpm at 29 °C. After 24 h, mycelium was transferred to aflatoxin inducing medium, YES, for 6 and 24 h. Each strain was grown three times.

3.2. *LaeA* is required for aflatoxin production, *aflR* expression and normal *veA* expression

Drawing from observations of *A. fumigatus* and *A. nidulans* $\Delta laeA$ strains, we predicted a loss of aflatoxin production mediated by loss of expression of the aflatoxin specific transcription factor *aflR* in the *A. flavus* $\Delta laeA$ strain. As shown in Fig. 1A and Supplementary Fig. 2, while the $\Delta laeA$ mutants did not show any detectable aflatoxin production, aflatoxin production was restored and in fact increased in TJW79.13 (and other complemented strains) as compared to wildtype. Examination of *laeA* and *aflR* expression in wildtype, TJW71.1 and TJW79.13 showed no detectable expression of either *aflR* or *laeA* in the $\Delta laeA$ mutant but increased expression of both genes in TJW79.13 (Fig. 1B), thus correlating with TLC results.

In addition to *aflR* expression, we also examined expression of *veA*. VeA is a light regulated gene with similar global effects on secondary metabolite production in all the examined aspergilli (Stinnett et al., 2007). VeA is necessary for aflatoxin production and *aflR* expression in *A. parasiticus* (Calvo et al., 2004) and another isolate of *A. flavus* (Duran et al., 2007). Most recently VeA and LaeA, along with a third protein VeIB, have been shown to function as a nuclear complex in *A. nidulans* (Bayram et al., 2008). As shown in Fig. 1B, in comparison to wildtype, the *laeA* deletion mutant TJW71.1 showed an increase and TJW79.13 a decrease in *veA* expression. These results suggest that *veA* is negatively regulated by LaeA.

3.3. Decreased conidial production and loss of sclerotia in $\Delta laeA$

Similarly to *A. fumigatus* and *A. nidulans* $\Delta laeA$ strains, the *A. flavus* $\Delta laeA$ mutants presented a loss of pigmentation on the backside of GMM growth plates, although no difference in radial growth (data not shown). Quantitative analysis of conidial production showed an approximate two-fold decrease in spore production in the $\Delta laeA$ mutant with no observable phenotype in the

OE::laeA strain on GMM medium (Fig. 2A). This decrease was also observed in YES medium (Fig. 2B).

Sclerotial production has been linked to aflatoxin synthesis in several studies including the loss of both processes in ΔveA (Calvo et al., 2004; Duran et al., 2007). To determine if loss or overexpression of *laeA* resulted in aberrations in sclerotial production, the strains were grown on sclerotial inducing medium (GMM medium supplemented with 2% sorbitol). Striking phenotypes were observed for both mutants where the $\Delta laeA$ strain did not produce any sclerotia and the OE::laeA strain produced statistically increased sclerotial numbers compared to wildtype (Fig. 3).

3.4. *laeA* mutants are altered in host colonization

Although aflatoxin is not considered a virulence factor and alterations in its production would not a priori be considered important in seed colonization, we considered it possible that other changes wrought by deletion or overexpression of *laeA* could affect colonization of host seed. To examine this possibility, peanut seeds were inoculated with all three strains of *A. flavus*.

Macroscopically, the $\Delta laeA$ strain appeared to grow less vigorously than wildtype on peanut, whereas the OE::laeA strain showed an early enhanced ability to colonize peanut seeds (Fig. 4A), however, by day three any macroscopic difference was largely obscured (data not shown). Visual observations were correlated with conidial production by the OE::laeA strain on seed. On both day 2 and 3, the OE::laeA strain produced statistically more conidia than both wildtype and $\Delta laeA$ (Fig. 4B and data not shown). Although there was no decrease in conidial production by $\Delta laeA$ on peanut seed as assessed by these assays, there was on maize seed (Supplementary Fig. 3).

Fig. 4C shows that aflatoxin production on the peanut seed reflected that of production on medium (Fig. 1A), namely that $\Delta laeA$ did not produce aflatoxin on seed whereas the OE::laeA produced more than wildtype. A similar aflatoxin profile was seen on maize




Fig. 2. *LaeA* deletion results in reduced sporulation on growth medium. (A) The wildtype *A. flavus* NRRL 3357 and three $\Delta laeA$ mutants (TJW71.1, TJW71.7 and TJW71.3) were grown in YES medium for 5 days at 29 °C. Columns with the same letter are not significantly different at a significance level of 0.05. (B) The wildtype *A. flavus* NRRL 3357 and three $\Delta laeA$ mutants (TJW71.1, TJW71.7 and TJW71.3) were grown on GMM medium for 5 days. Columns with the same letter are not significantly different at a significance level of 0.01. (C) The wildtype NRRL 3357, $\Delta laeA$ (TJW71.1) and *OE::laeA* (TJW79.13) where grown on GMM medium at 29 °C for 5 days. Columns with the same letter are not significantly different at a significance level of 0.05. For all tests, conidia were counted using a hemocytometer. Values are a mean of four replicates.

seed (where only $\Delta laeA$ strains were examined, Supplementary Fig. 3).

Several enzyme activities have been associated with *Aspergillus* pathogenesis of seed, including lipase activity (Tsitsigiannis and Keller, 2006). We assessed potential lipase activity for all three strains. As shown in Fig. 5, $\Delta laeA$ showed lower lipase activity than either wildtype or the *OE::laeA* strain at both day 2 and 4.

3.5. Global regulation of secondary metabolites in *LaeA* mutants

LaeA regulates production of several secondary metabolites in *A. fumigatus* and *A. nidulans* (Bok and Keller, 2004; Perrin et al., 2007). Here we examined the production of known compounds produced by wildtype *A. flavus* NRRL 3357 (Frisvad et al., 1987)




Fig. 3. Sclerotial production in $\Delta laeA$ and *OE::laeA* mutants. The wildtype NRRL 3357, $\Delta laeA$ (TJW71.1) and *OE::laeA* (TJW79.13) where grown on GMM plus 2% sorbitol medium at 29 °C for 5 days in the dark. Plates were sprayed with ethanol to wash off conidia to visualize sclerotia. Sclerotia were absent in the $\Delta laeA$ strain and overproduced in the *OE::laeA* strain as compared to wildtype. The average number of sclerotia produced per petri plate by the *OE::laeA* strain (1540 ± 90) was statistically greater ($P < 0.01$) than the average number produced by wildtype (217 ± 70).

in comparison to both *laeA* mutants. The profile of these secondary metabolites depended on the growth substrate (Table 1). Two striking observations were made. First, the $\Delta laeA$ mutant did not produce any secondary metabolites, with the exception of kojic acid on DG18 agar and aspergillilic acid on TGY agar. Second, the *OE::laeA* strain produced additional metabolites not observed in wildtype. These were all known sclerotial metabolites including paspaline/paspalicine, aflatrem and aflavinines and their production was attributed to the high production of sclerotia in this strain. The relative amounts of aflatoxins were slightly higher in the *OE::laeA* strain than in the wildtype on CYA and DG18 agar, while the opposite was the case for YES agar. The amount of kojic acid was 3–10 times higher for the wildtype than the *OE::laeA* strain on YES agar, but the opposite was the case for DG18 agar.

4. Discussion

A critical attribute in *LaeA* function is its role in transcriptional regulation of *Aspergillus* secondary metabolite gene clusters. Genes required for expression of these biologically active small molecules are typically arranged in a contiguous manner and inherited as a single genetic locus (Hoffmeister and Keller, 2007). *LaeA* activity allows for directed expression of many of these loci with loss of *laeA* resulting in gene silencing (Bok et al., 2006b; Perrin et al., 2007). Although *LaeA* cellular mechanism is yet unknown, accruing data suggests *LaeA* may play a role in epigenetic regulation of such clusters (Bok et al., 2006c; Shwab et al., 2007), possibly as a member of a nuclear complex with VeA (Bayram et al., 2008). *LaeA* governs expression of multiple gene clusters in the soil saprophyte *A. nidulans* (Bok et al., 2006b). Feeding choice preferences of a fungivore demonstrated that *laeA* provides both a protective shield for the fungus as well as negatively impacts egg laying in the consuming springtails (Rohlfes et al., 2007). Similarly, *LaeA* function is crit-




Fig. 4. Growth and aflatoxin production on peanut seed. The wildtype NRRL 3357, $\Delta laeA$ (TJW71.1) and $OE::laeA$ (TJW79.13) were grown on peanut seed for 2 days at 29 °C. (A) Representative picture of inoculated seed. (B) Conidia were counted from four replicates of seed as pictured in (A). Columns with the same letter are not significantly different at a significance level of 0.05. (C) Aflatoxin production on peanut seed as examined by thin layer chromatography. C is control (seed mock inoculated with water). AF is aflatoxin B1 standard.




Fig. 5. Lipase activity in $\Delta laeA$ and $OE::laeA$ mutants. The wildtype NRRL 3357, $\Delta laeA$ (TJW71.1) and $OE::laeA$ (TJW79.13) were grown on lipase assay medium where increased clearing zone indicates increased lipase-like activity. Values are a mean of four replicates. Columns with the same letter are not significantly different at a significance level of 0.05. Day 2 and day 4 were analyzed separately.

ical for production of several toxins in *A. fumigatus*, including gliotoxin, a molecule injurious to mammalian cells and possibly important in the severity of IA as determined by the animal models and cellular response (Bok et al., 2006a; Spikes et al., 2008; Sugui et al., 2007b). These studies have strongly implicated loss of metabolite production as a key factor contributing to the reduced virulence of $\Delta laeA$ strains (Bok et al., 2005; Sugui et al., 2007a).

To address an impact on *LaeA* in *A. flavus* physiology and pathogenicity, we examined both $\Delta laeA$ and $OE::laeA$ strains. Our analysis showed reduced (no) expression of cyclopiazonic acid, kojic acid, oryzaechlorin and asperfurin in addition to aflatoxin in the $\Delta laeA$ mutant, whereas several metabolites were up regulated in the $OE::laeA$ strain, particularly those associated with sclerotial production, e.g. paspaline, aflatrem and aflavinines (Table 1). Although not produced by $\Delta laeA$ in most media, kojic acid was produced when the strain was grown on DG18 medium. The production of aspergillic acid was approximately the same in all three isolates and appears not to be controlled by *laeA*. Both aspergillic acid and kojic acid are putative chelation agents and, as such, possibly play an essential role in general fitness of the fungus, which may impact the degree of regulation by *LaeA* of these metabolites. Interestingly, one metabolite, oryzaechlorin, an epidithiodiketopiperazine similar in structure to gliotoxin, was down regulated in both mutants.

Loss of metabolite production has been correlated with morphological differentiation in fungi (Calvo et al., 2002), including reduced conidiospore formation (Wilkinson et al., 2004) and sclerotial loss (Calvo et al., 2004) in various *Aspergilli*. Although aflatoxin is not correlated with disease severity, sclerotial production has been associated with pathogenicity in some *Aspergilli* including *A. flavus* (Chang et al., 2001). The loss and increased sclerotial production of the two *laeA* mutants, and associated metabolites, was striking (Fig. 3 and Table 1) and reminiscent of the *AveA* phenotype. Deletion of *veA* in *A. parasiticus* (Calvo et al., 2004) and another strain of *A. flavus* (Duran et al., 2007) also show loss of *aflR* expression (and consequently aflatoxin production) coupled with

Table 1Production of secondary metabolites by wildtype (NRRL 3357), *ΔlaeA* (TJW71.1) and *OE::laeA* (TJW79.13) as measured by HPLC-DAD (diode array detection)

<i>A. flavus</i> strains	Aflatoxin B ₁ RI 835	Aflatoxin B ₂ RI 810	Cyclopiazonic Acid RI 1094	Kojic Acid RI 570	Aspergillitic Acid RI 1976	Paspaline/paspalinine RI 1261/1479	Aflatrems RI 1522	Aflavinines ^a	Oryzochlorin RI 850
3357 CYA	2.5 34 LU ^b	ND 32 LU	2.8	0	0	0	0	0	0
<i>ΔlaeA</i> CYA	0	ND	0	0	0	0	0	0	0
<i>OE::laeA</i> CYA	2.8 43 LU	ND 41 LU	2.8	0	0	1.8/2.0	1.9	2.3/0/1.6/2.0/2.0	0
3357 YES	3.1 84 LU	ND 102 LU	2.5	4.0	0	0	0	0	0
<i>ΔlaeA</i> YES	0	0	0	0	0	0	0	0	0
<i>OE::laeA</i> YES	2.9 58 LU	ND 90 LU	2.4	3.0	0	0	0	0	0
3357 WATM	2.0 26 LU	ND 27 LU	2.7	0	0	0	0	0	0
<i>ΔlaeA</i> WATM	0	0	0	0	0	0	0	0	0
<i>OE::laeA</i> WATM	2.5 26 LU	ND 28 LU	2.5	0	0	0/1.8	1.6	0/1.6/2.0/1.9/1.9	0
3357 TGY	0	0	0	0	2.1	0	0	0	0
<i>ΔlaeA</i> TGY	0	0	0	0	2.0	0	0	0	0
<i>OE::laeA</i> TGY	0	0	0	0	1.9	0	0	0	0
3357 DG18	1.6 LU 2	ND LU 0	0	4.0	0	0	0	0	1.9
<i>ΔlaeA</i> DG18	0	0	0	3.3	0	0	0	0	0
<i>OE::laeA</i> DG18	2.6 LU 29	ND LU 37	1.9	3.5	0	0	0	0	0

The log to the area of the peaks is given as mAU (milli absorption units) as measured at 210 nm (ND means non detectable at 210 nm).

^a The aflavinines had retention indices (RI) 967, 1146, 1185, 1610, 1650.

^b Aflatoxin B₁ and B₂ as measured by a fluorescence detector (LU units, excitation 230 nm, emission 450 nm).

loss of sclerotial formation. Examination of *veA* expression in *ΔlaeA* and *OE::laeA* strains supports a LaeA mediated negative regulatory mechanism governing *veA* expression. A coupling of all studies would suggest that both *veA* and *laeA* are required for aflatoxin and sclerotial formation, perhaps through formation of their joint regulatory complex in the nucleus (Bayram et al., 2008). Possibly regulation of *veA* by LaeA represents an internal mechanism to balance stoichiometry of this complex.

We also noted, based on visual observations and softness of infected seed, that *ΔlaeA* appeared less aggressive in tissue maceration compared to wildtype or the *OE::laeA* strain (Fig. 4A). As one indicator of maceration potential, we assessed overall lipase/esterase activity of these mutants, a technique successfully employed in other studies to assess fungal degradative potential (Tsitsigiannis and Keller, 2006). Degradative enzymes are well known to play a significant role in fungal infections of plants, with lipases particularly important in pathogenesis of several hosts including seed in the *A. flavus* pathosystem (Berto et al., 1999; Commenil et al., 1995). The decreased lipase activity of the *ΔlaeA* mutant may indicate lipase activity contributes to pathogenesis through tissue degradation in this strain (Fig. 5). The differences in conidial production of this mutant on peanut and maize seed may reflect host differences or could be an attribute of timing of assessment and/or method of inoculation.

In summary, results from this study demonstrate the conserved function of *laeA* as a global regulator of secondary metabolism in the Aspergilli with varying impact on *Aspergillus* morphology. We speculate the development of chemical regulation by LaeA may have evolved as a protective device (Rohlfes et al., 2007) and, by chance, also provides considerable aggressive characteristics in disease development by opportunistic fungi. Regardless of the role of LaeA in fungal biology, our findings represent an advance in identification of a shared mechanism in *Aspergillus* resulting in damage of both plant and animal hosts.

5. Seed infections

5.1. Peanut (*Arachis hypogaea*)

Mature peanut seeds were prepared by removing the brown exterior peanut layer (testa) using the fingers. The two cotyledons of each seed were separated and the embryo carefully removed

without damaging any of the cotyledon tissue. Then, cotyledons were surface sterilized by placing them in a tea ball infuser and dipping them in a beaker containing 0.05% sodium hypochlorite in sterile water for 3 min. The tea ball was transferred to a new beaker containing sterile distilled water for 30 s (wash step), followed by a 5-s wash with 70% ethanol in a new beaker (additional sterilization step) and one more 30-s wash with sterile distilled water while shaking the tea ball. The cotyledons were drained completely and placed in a petri dish until the time of infection. All the steps were aseptically performed in a biosafety hood.

Peanut cotyledons were inoculated with a 10⁵ spores/ml. Cotyledon treatments included water control (mock inoculation) and infection with fungal strains. For all treatments, 20 peanut cotyledons were immersed in 20 ml of sterile distilled water (control) or sterile distilled water with fungal conidia in 50 ml centrifuge tubes while shaking for 30 min in a rotary shaker at 50 rpm. Cotyledons were placed in petri dishes lined with three pieces of moist filter paper (to create a humidity chamber) and a water reservoir (lid of a 50 ml centrifuge tube containing 2 ml of sterile water) to maintain high humidity. Cotyledons were incubated for 3 days at 29 °C in dark conditions. The filter paper was moistened daily.

5.2. Maize (*Zea mays*)

Untreated maize seeds (X516WX raw) obtained from Kaltenberg Seed Farms, Inc. (Waunakee, WI) were inoculated similarly to peanut seed with the following differences. Seeds were wounded by creating a small hole near the embryo where they were inoculated with 5 μl of a 10⁶ spores/ml suspension of *A. flavus* conidia. Ten seeds per replicate were transferred to the humidity chamber and incubated in the dark at 29 °C for 3 days. The filter paper was moistened daily and any germinating seeds were removed and discarded. Analysis was performed on the same number of seed for each treatment and replication (e.g. if three seed germinated from one humidity chamber, then only seven seed were examined for each treatment and replication).

All seed experiments were repeated three to four times.

5.3. Aflatoxin analysis from seed

Three days after infection, peanut cotyledons were collected in 50 ml centrifuge tubes with the addition of 3 ml of 0.01% Tween 80

(v/v in water) and vortexed vigorously for 1 min. One milliliter was removed from each sample for conidia counting prior to aflatoxin extraction. For extraction, 5 ml of acetone was added to the samples followed by shaking for 10 min in a rotary shaker at 150 rpm. Samples were allowed to stand for 5 min at room temperature and then 5 ml of chloroform was added to each sample followed by shaking for 10 min at 150 rpm. Samples were allowed to stand for an additional 10 min at room temperature, vortexed briefly and centrifuged for 15 min at 2000 rpm to collect the organic lower phase. Samples were then dried out completely. The presence of abundant seed lipids in the samples hampered the clear observation of aflatoxin on TLC plates and a second extraction-purification was carried out as follows. Samples were re-suspended in 5 ml of 0.1 M NaCl methanol:water (55:45) and 2.5 ml of hexane and vortexed vigorously at high speed for 1 min. Samples were centrifuged at 2000 rpm for 5 min. The hexane layer was collected and the fatty acid inter-phase layer was discarded. The remaining aqueous phase was washed an additional time with 2.5 ml hexane as described above. The hexane extracts were combined, allowed to dry and then re-suspended in 500 μ l of chloroform before 10 μ l of each extract was separated on a silica gel TLC plate using a chloroform:acetone (95:5 [vol/vol]) solvent system.

Maize seed were extracted similarly but without the needed extra hexane extraction for peanut seed.

5.4. Lipase assays

To test for lipase activity, GMM medium was overlaid with 10 ml of a 10^6 spores/ml suspension of *A. flavus* conidia in molten agar. Cultures were grown for 4 days at 29 °C in light conditions. Then, one 0.7-cm diameter agar plug was harvested from the center of each plate and added to a sterile test tube containing lipase medium (0.5% mycological peptone, 0.3% yeast extract in 1% agar containing 0.1% glyceryl tributyrates, Paterson et al., 1994). Tubes were incubated at 29 °C in light conditions. Measurements of the clearing zone, indicative of lipase activity, were taken at days 2 and 4.

5.5. Aflatoxin production in media

YES and GMM media were overlaid with 5 ml of a 10^6 spores/ml suspension of *A. flavus* conidia in molten agar. Cultures were grown for 5 days at 29 °C. Three 1.5-cm diameter cores were harvested from the center of each plate and homogenized in 3 ml of distilled water. Three milliliter each of acetone and chloroform were added and the mixture was vortexed for 1 min. The mixture was incubated at room temperature for 1 h, was vortexed again and centrifuged for 10 min. The lower organic layer was removed and evaporated. Residue was resuspended in 100 μ l of chloroform and 10 μ l of the suspension were spotted onto a TLC plate (Whatman Ltd., Kent, England). This experiment was done in triplicate. Aflatoxin was resolved in a chloroform and acetone (95:5) solvent system and standards were purchased from Sigma-Aldrich (St. Louis, MO). Aflatoxin was visualized using long-wave (366 nm) UV light and digital photographs were taken.

5.6. Comprehensive secondary metabolite profiles

Three isolates *A. flavus* NRRL 3357, *A. flavus* TJW71.1 (Δ laeA) and *A. flavus* TJW79.13 (OE::laeA) were inoculated on the media CYA, YES (Frisvad and Samson, 2004), DG18 (Dichloran 18% glycerol agar, http://www.cbs.knaw.nl/cbs_home/cbs_home.html?http://www.cbs.knaw.nl/food/media.htm-main), WATM (Wickerhams Antibiotic test medium, (Raper and Thom, 1949)), and TGY (tryptone glucose yeast extract agar, <http://www.cbs.knaw.nl/>

[cbs_home/cbs_home.html?http://www.cbs.knaw.nl/food/media.htm-main](http://www.cbs.knaw.nl/food/media.htm-main)) (in three points). The media were incubated for 7 days in the dark and five agar plugs were taken for HPLC analysis. The agar plugs were extracted with ethyl acetate/dichloromethane/methanol (3:2:1) with 1% formic acid added to this mixture and the extract ultrasonicated for 50 min. The extract was transferred to a new vial, and the organic solvents evaporated. The dry extract was re-dissolved in 0.5 ml methanol and filtered through a 0.45 μ m filter before analysis. 3 μ l of extract were injected into the chromatograph. The method used was based on Smedsgaard (Smedsgaard, 1997), but a Luna C18 (II) (Phenomenex, USA) column was used and the running time was 25 min. Retention indices were calculated for each compound and the retention indices and UV spectra were compared with authentic standards (Frisvad and Thrane, 1987).

5.7. Northern blot analysis of laeA, aflR and veA

Fifty milliliters of liquid YEP medium (6% peptone, 2% yeast extract) was inoculated with 10^6 conidia/ml of *A. flavus* NRRL 3357, TJW71.1, or TJW79.13 in 50 ml flasks, incubated with shaking at 250 rpm at 29 °C. After 24 h, the mycelium was collected and incubated in the aflatoxin-stimulating YES medium for 6 and 24 h (220 rpm, 29 °C). Mycelia were harvested and total RNA was isolated using the Trizol (source: Invitrogen) method. The Northern blot was hybridized with a 670 bp HindIII/BamHI *aflR* fragment from pTMH52.1 (McDonald et al., 2005), a 4.3 kb *laeA* fragment from pLRM11 and a 1.7 kb *veA* fragment generated by PCR using forward primer 5' CTAGCTGGTCATTATTTGATCTCG and reverse primer 5' GTTGTAGAGTGGACGATCATCATG from genomic DNA.

5.8. Statistical analysis

Statistical differences in spore numbers were calculated by ANOVA tests using Minitab software (Penn State University).

Acknowledgments

Our thanks to Dr. Gary A. Payne for access to *A. flavus* genome data and for providing us with strains. This research was funded by NIH (MBRS) Grant 5SO6GM08008 to S.P.K., and in part by NSF MCB-0236393, USAID Peanut Collaborative Research Support Program Grant LAG-G-00-96-90013-00 and USDA/NRI Competitive Grant 2002-35201-12562 to N.P.K. and NIH R01 AI065728-01A1 to N.P.K. and J.C.F. J.C.F. also thanks the Danish Research Agency for financial support (Center for Microbial Biotechnology).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.fgb.2008.06.009.

References

- Bayram, Ö. et al., 2008. VelB/VeA/LaeA coordinate light signal with fungal development and secondary metabolism. *Science* 320, 1504–1506.
- Berto, P. et al., 1999. Occurrence of a lipase in spores of *Alternaria brassicicola* with a crucial role in the infection of cauliflower leaves. *FEMS Microbiol. Lett.* 180, 183–189.
- Bok, J.W. et al., 2005. LaeA, a regulator of morphogenetic fungal virulence factors. *Eukaryot. Cell* 4, 1574–1582.
- Bok, J.W. et al., 2006a. GliZ, a transcriptional regulator of gliotoxin biosynthesis, contributes to *Aspergillus fumigatus* virulence. *Infect. Immun.* 74, 6761–6768.
- Bok, J.W. et al., 2006b. Genomic mining for *Aspergillus* natural products. *Chem. Biol.* 13, 31–37.
- Bok, J.W., Keller, N.P., 2004. LaeA, a regulator of secondary metabolism in *Aspergillus* spp. *Eukaryot. Cell* 3, 527–535.
- Bok, J.W. et al., 2006c. Secondary metabolic gene cluster silencing in *Aspergillus nidulans*. *Mol. Microbiol.* 61, 1636–1645.

- Bressac, B. et al., 1991. Selective G to T mutations of p53 gene in hepatocellular carcinoma from southern Africa. *Nature* 350, 429–431.
- Calvo, A.M. et al., 2004. *veA* is required for toxin and sclerotial production in *Aspergillus parasiticus*. *Appl. Environ. Microbiol.* 70, 4733–4739.
- Calvo, A.M. et al., 2002. Relationship between secondary metabolism and fungal development. *Microbiol. Mol. Biol. Rev.* 66, 447–459.
- Chang, P.K. et al., 2001. Association of aflatoxin biosynthesis and sclerotial development in *Aspergillus parasiticus*. *Mycopathologia* 153, 41–48.
- Commenil, P. et al., 1995. Purification and properties of an extracellular lipase from the fungus *Botrytis cinerea*. *Lipids* 30, 351–356.
- Cove, D.J., 1966. The induction and repression of nitrate reductase in the fungus *Aspergillus nidulans*. *Biochim. Biophys. Acta* 113, 51–56.
- Duran, R.M. et al., 2007. Production of cyclopiazonic acid, aflatrem, and aflatoxin by *Aspergillus flavus* is regulated by *veA*, a gene necessary for sclerotial formation. *Appl. Microbiol. Biotechnol.* 73, 1158–1168.
- Frisvad, J.C., Samson, R.A., 2004. Polyphasic taxonomy of *Penicillium* subgenus *Penicillium*—A guide to identification of food and air-borne terverticillate *Penicillia* and their mycotoxins. *Stud. Mycol.*, 1–173.
- Frisvad, J.C., Thrane, U., 1987. Standardized high-performance liquid chromatography of 182 mycotoxins and other fungal metabolites based on alkylphenone retention indices and UV-vis spectra (diode array detection). *J. Chromatogr.* 404, 195–214.
- Hedayati, M.T. et al., 2007. *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology* 153, 1677–1692.
- Hoffmeister, D., Keller, N.P., 2007. Natural products of filamentous fungi: enzymes, genes, and their regulation. *Nat. Prod. Rep.* 24, 393–416.
- Latge, J.P., 1999. *Aspergillus fumigatus* and aspergillosis. *Clin. Microbiol. Rev.* 12, 310–350.
- Liang, S.H. et al., 1997. Analysis of mechanisms regulating expression of the *ver-1* gene, involved in aflatoxin biosynthesis. *Appl. Environ. Microbiol.* 63, 1058–1065.
- McDonald, T. et al., 2005. Inverted repeat transgenes silence mycotoxin production in *Aspergillus* and *Fusarium* species. *Mol. Plant Microbe Interact.* 18, 539–545.
- Paterson, R.R.M. et al., 1994. *Biochemical Techniques for Filamentous Fungi*. CAB International, Wallingford.
- Payne, G.A., Brown, M.P., 1998. Genetics and physiology of aflatoxin biosynthesis. *Annu. Rev. Phytopathol.* 36, 329–362.
- Perrin, R.M. et al., 2007. Transcriptional regulation of chemical diversity in *Aspergillus fumigatus* by *LaeA*. *PLoS Pathog.* 3, e50.
- Probst, C. et al., 2007. Outbreak of an acute aflatoxicosis in Kenya in 2004: identification of the causal agent. *Appl. Environ. Microbiol.* 73, 2762–2764.
- Raper, K.B., Thom, C., 1949. *A Manual of the Penicillia*. Williams & Wilkins Co., Baltimore.
- Richard, J.L., Payne, G.A., 2003. Mycotoxins: risks in plant, animal, and human systems. Task Force Report No. 139. Council for Agricultural Science and Technology, Ames, IA, pp. xvi+ 199.
- Rohlf, M. et al., 2007. Secondary chemicals protect mould from fungivory. *Biol. Lett.* 3, 523–525.
- Samson, R.A., 1992. Current taxonomic schemes of the genus *Aspergillus* and its teleomorphs. In: Klich, J.W.B.M.A. (Ed.), *Aspergillus: The Biology and Industrial Applications*. Butterworth Publishers, pp. 353–388.
- Shimizu, K., Keller, N.P., 2001. Genetic involvement of a cAMP-dependent protein kinase in a G protein signaling pathway regulating morphological and chemical transitions in *Aspergillus nidulans*. *Genetics* 157, 591–600.
- Shwab, E. et al., 2007. Histone deacetylase activity regulates chemical diversity in *Aspergillus*. *Eukaryot. Cell* 6, 1656–1664.
- Smedsgaard, J., 1997. Micro-scale extraction procedure for standardized screening of fungal metabolite production in cultures. *J. Chromatogr. A* 760, 264–270.
- Spikes, S. et al., 2008. Gliotoxin production in *Aspergillus fumigatus* contributes to host-specific differences in virulence. *J. Infect. Dis.* 197, 479–486.
- Stinnett, S.M. et al., 2007. *Aspergillus nidulans* *VeA* subcellular localization is dependent on the importin alpha carrier and on light. *Mol. Microbiol.* 63, 242–255.
- Sugui, J.A. et al., 2007a. Role of *laeA* in the regulation of *alb1*, *gliP*, conidial morphology, and virulence in *Aspergillus fumigatus*. *Eukaryot. Cell* 6, 1552–1561.
- Sugui, J.A. et al., 2007b. Gliotoxin is a virulence factor of *Aspergillus fumigatus*: *gliP* deletion attenuates virulence in mice immunosuppressed with hydrocortisone. *Eukaryot. Cell* 6, 1562–1569.
- Tsitsigiannis, D.I., Keller, N.P., 2006. Oxylipins act as determinants of natural product biosynthesis and seed colonization in *Aspergillus nidulans*. *Mol. Microbiol.* 59, 882–892.
- Wilkinson, H.H. et al., 2004. Increased conidiation associated with progression along the sterigmatocystin biosynthetic pathway. *Mycologia* 96, 1190–1198.