

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/303699299>

New natural products from the marine-derived *Aspergillus* fungi–A review

Article in ACTA MICROBIOLOGICA SINICA · December 2015

DOI: 10.13343/j.cnki.wsxb.20150478

CITATIONS

0

READS

25

1 author:



Weiming Zhu

Ocean University of China

214 PUBLICATIONS 2,480 CITATIONS

SEE PROFILE



海洋曲霉来源的新天然产物

赵成英, 刘海珊, 朱伟明*

中国海洋大学医药学院, 海洋药物教育部重点实验室, 山东 青岛 266003

摘要: 海洋真菌由于其遗传背景复杂、代谢产物种类多且产量高, 已成为海洋微生物新天然产物的主要来源, 从我们对2010–2013年初的海洋微生物来源新天然产物的统计来看, 研究最多的是曲霉属(*Aspergillus*)真菌, 占海洋真菌来源新天然产物的31%。本文从菌株来源、化合物结构及其生物活性等方面, 综述了自1992年第一个海洋曲霉天然产物到2014年8月已报道的共512个海洋曲霉来源的新天然产物。这些海洋天然产物具有丰富的化学多样性, 且36%的化合物表现出细胞毒、抑菌、抗氧化和抗寄生虫等生物活性; 含氮化合物是主要的结构类型、约占曲霉源海洋天然产物总数的52%, 也是出现活性化合物比例最高的结构类型、约40%的含氮化合物具有生物活性, 其中脱氢二酮哌嗪生物碱halimide的化学衍生物plinabulin已结束II期临床研究, 并于2015年第三季度开始在美国和中国进行III期临床研究, 用于治疗转移性的晚期非小细胞肺癌。

关键词: 海洋真菌, 曲霉, 天然产物, 化学结构, 生物活性, 来源

海洋真菌由于其遗传背景复杂、代谢产物种类多、产量高, 成为海洋微生物新天然产物的主要来源, 作者对2010–2013年初的海洋微生物来源新天然产物的统计表明, 研究最多的真菌属是曲霉(*Aspergillus*), 占海洋真菌新天然产物的31%^[1]。海洋曲霉源天然产物的研究始于1992年, Shinggu等报道了首例海洋曲霉来源的新天然产物fumiquinazolines A–C (**149–151**)^[2]; 截止到2014年8月, 已报道512个海洋曲霉来源的新天然产物。其结构类型多样(包括聚酮、生物碱、萜类、甾

体、卤代物、脂肪酸、肽类、糖苷等), 且有多种生物活性(包括抗癌、抑菌、自由基清除和抗寄生虫等)。值得一提的是, 由海洋曲霉天然产物halimide(**301**)^[3–4]衍生而来的plinabulin (NPI-2358)^[5]是一种血管阻断剂, 作用于肿瘤细胞, 影响微管蛋白解聚; 目前plinabulin已结束II期临床研究、开始在美国和中国进行III期临床研究, 用于治疗转移性的晚期非小细胞肺癌(NSCLC)^[6], 成为20个海洋药物之一, 也是唯一的海洋曲霉菌来源的药物^[7]。由此可见海洋曲霉是海洋天然产物

基金项目: 国家自然科学基金(41376148, 81561148012, U1406402-1, U1501221)

*通信作者: Tel/Fax: +86-532-82031836; E-mail: weimingzhu@ouc.edu.cn

收稿日期: 2015-10-15; 修回日期: 2015-12-25; 网络出版日期: 2015-12-30

乃至新药发现的重要资源, 本文将综述这些海洋曲霉新天然产物的菌株来源、结构及其生物活性。

1 海洋动物来源的曲霉天然产物

1.1 海绵来源的曲霉天然产物

氯代物chlorocarolides A(**1**)和B(**2**)来源于赫曲霉*Aspergillus cf. ochraceus* 941026^[8]。1株黑曲霉*Aspergillus niger*代谢产生二酮哌嗪(环缩二氨酸)的二聚体asperazine(**3**), 该化合物可以选择性抑制人白血病细胞L1210、C38以及人结肠癌细胞H116或者CX1细胞株^[9]。化合物asperic acid(**4**)是从另外1株黑曲霉*A. niger* 94-1212的次生代谢产物中分离得到^[10]。花斑曲霉*A. versicolor* (Vuill) Triab代谢产生六元色酮的衍生物aspergiones A-F(**5**–**10**)^[11]以及aspergillone(**11**)、aspergillodiol(**12**)、aspergillol(**13**)和12-acetyl-aspergillol(**14**)^[12]。3个含氯的抗生素**15**–**17**来自孔曲霉*A. ostianus* TUF 01F313, 均对大西洋鲁杰氏菌*Ruegeria atlantica*有抑制活性, 其中化合物**16**和**17**在25 $\mu\text{g}/\text{disc}$ 浓度下的抑菌圈直径分别为10.1 mm和10.5 mm, 而化合物**15**在5 $\mu\text{g}/\text{disc}$ 时即表现出较好的抑制活性(抑菌圈直径为12.7 mm), 另外该化合物还有微弱的金黄色葡萄球菌抑制活性(25 $\mu\text{g}/\text{disc}$ 时抑菌圈直径为10.2 mm)^[13]。进一步研究还分离到aspinotriols A(**18**)和B(**19**)、aspinonediol(**20**)^[14], aspergillides A–C(**22**–**24**)^[15]及其七环生物碱化合物21-hydroxystephacidin A(**25**)^[16](图1)。并确定了dihydroaspyrone(**21**)的绝对构型^[14], 其中化合物**22**–**24**对L1210细胞的 LD_{50} 分别为2.1、71.0和2.0 $\mu\text{g}/\text{mL}$ ^[15]。

Circumdatins D–F(**26**–**28**)分离自另外一株孔曲霉*A. ostianus* IBT 12704^[17]。混源萜tropolactones A–D(**29**–**32**)由曲霉属*Aspergillus* sp. CNK-371代谢产生, 其中化合物**29**–**31**对HCT-116细胞有微弱的抑制活性, IC_{50} 分别为13.2、10.9和13.9 $\mu\text{g}/\text{mL}$ ^[18]。

棘孢曲霉*A. aculeatus* CRI323-04代谢产生aspergillusol A(**33**)^[19]和asperaculin A(**34**)^[20], 其中化合物**33**对酵母和嗜热脂肪芽孢杆菌来源的 α -糖苷酶有抑制活性, IC_{50} 值分别为465和1060 $\mu\text{mol}/\text{L}$ ^[19]。化合物**35**–**37**来自另一株棘孢曲霉*A. aculeatus* CRI322-03^[21]。曲霉*A. insuetus*代谢产生Terretonins E(**38**)和F(**39**), 均具有哺乳动物线粒体呼吸链抑制作用, IC_{50} 值分别为3.90和2.97 $\mu\text{mol}/\text{L}$ ^[22]。菌核曲霉*A. sclerotiorum* Huber SP080903f04代谢产生N-去甲棕曲菌素JBIR-15(**40**)^[23]。焦曲霉*A. ustus* 8009代谢产生7个补身烷倍半萜化合物**41**–**47**, 其中**44**、**45**对多种肿瘤细胞有细胞毒活性, 尤其是**45**对L5178Y细胞株的 EC_{50} 值为0.6 $\mu\text{g}/\text{mL}$ ^[24]。蛇孢甲壳素**48**–**52**及吡咯生物碱**53**、**54**也来自同一菌株^[25]。脂肽fellutamide C(**55**)(图1)来自于*A. versicolor*, 对SK-MEL-2、XF498和HCT15的 IC_{50} 分别为5.1、3.9和3.1 $\mu\text{mol}/\text{L}$ ^[26]。

JBIR-74(**56**)和JBIR-75(**57**)来自*Aspergillus* sp. fs14^[27]。混源萜insuetolides A–C(**58**–**60**)和补身烷倍半萜**61**来自奇突曲霉*A. insuetus* OY-207, 其中化合物**58**有抑制粗糙链孢霉菌的活性、MIC为140 $\mu\text{mol}/\text{L}$, 化合物**60**和**61**对人MOLT-4细胞有细胞毒活性、50 mg/mL 时的抑制率分别为51%和55%^[28]。脂肽fellutamide F(**62**)来自*A. versicolor* PF10M, 对多株人癌细胞的 EC_{50} 为0.13–1.81 $\mu\text{g}/\text{mL}$ ^[29]。*Aspergillus* sp.代谢产生4个没药烷倍半萜aspergiterpenoid A(**63**)、(–)-sydonol(**64**)、(–)-sydonic acid(**65**)和化合物**66**, 其对金黄色葡萄球菌、枯草芽孢杆菌、蜡状芽孢杆菌等有抑菌活性(MIC为1.25–20.0 $\mu\text{mol}/\text{L}$)^[30]。没药烷倍半萜二聚体disydonols A–C(**67**–**69**)也来自同一菌株, 其中化合物**67**和**69**对HepG-2和Cashi细胞均有细胞毒活性(IC_{50} 分别为9.31/12.40 $\mu\text{g}/\text{mL}$ 和2.91/10.20 $\mu\text{g}/\text{mL}$)^[31]。缩酚酸环醚**70**–**72**、二芳基醚**73**及吡喃酮**74**来自爪甲曲霉*A. unguis* CRI282-03, 其中化合物**70**–**72**均有芳香酶抑制活性(IC_{50} 分别为2.2、




图 1. 化合物1-92的结构

Figure 1. Structures of compounds 1-92.

4.1和0.7 $\mu\text{mol/L}$), 且**70**和**71**在黄嘌呤氧化测试中表现出自由基清除活性(IC_{50} 分别为16.0和<15.6 $\mu\text{mol/L}$)^[32]。二酮哌嗪二聚体eurocristatine(**75**)产自冠突散囊菌*Eurotium cristatum* KUFC 7356^[33], *E. cristatum*是小冠曲霉*A. cristatellus*的有性型。1株*Aspergillus* sp.代谢产生austalides M-Q(**76-80**)^[34]、tryptoquivaline K (**81**)和 fumiquinazolines K-P (**82-87**)^[35]以及生物碱**88**和混源萜austalide R(**89**)^[36](图1)。其中化合物**81**在10 $\mu\text{g/mL}$ 浓度下对小鼠淋巴瘤细胞L5178Y的抑制率为23%^[35], 化合物**88**能

选择性抑制弧菌(*Vibrio* sp.)生长, 而**89**则广谱抗菌、其对*Vibrioharveyi*、*V. proteolyticus*、*V. carchariae*、*Shewanella putrefaciens*、*Roseobacter litoralis*、*Pseudoalteromonas elyakovii*、*P. irgensii*和*Halomonas aquamarina*均有抑制作用^[36]。

杂色曲霉*A. versicolor* MF359代谢产生3个柄曲霉素**90-92**(图1), 其中化合物**92**对金黄色葡萄球菌和枯草芽孢杆菌均有抑制活性, MIC分别为12.500 $\mu\text{g/mL}$ 和3.125 $\mu\text{g/mL}$ ^[37]。黑曲霉*A. niger*代谢产生3,3-bicoumarin bicoumanigrin(**93**)、

aspernigrins A和B(**94**、**95**)以及pyranonigrins A–D(**96**–**99**)(图2); 化合物**93**在浓度1–20 $\mu\text{g/mL}$ 时具有人肿瘤细胞毒活性, 化合物**94**、**95**在浓度为50 $\mu\text{g/mL}$ 有轻微的肿瘤细胞毒活性; 化合物**95**具有神经保护作用^[38]。Nafuredin(**100**)产自黑曲霉 *Aspergillus niger* FT-0554, 表现出猪蛔虫 *Ascarissuum* 延胡索酸还原酶(NFRD)抑制活性, 其 IC_{50} 为12 nmol/L^[39]。

1.2 珊瑚来源的曲霉天然产物

土曲霉 *A. terreus* HKI0499 代谢产生 aspernolides

A、B(**101**、**102**)^[40]。(+)–methyl sydowate(**103**)、7-deoxy-7,14-didehydroxydic acid(**104**)和7-deoxy-7,8-didehydroxydic acid(**105**)(图2)来自一株 *Aspergillus* sp., 其中化合物**103**可抑制金黄色葡萄球菌、100 $\mu\text{g/mL}$ 浓度下抑菌圈直径为11 mm^[41]。Aspergilones A、B(**106**、**107**)来自同一株菌, 其中化合物**106**对HL-60、MCF-7和A549细胞的 IC_{50} 分别为3.2、25.0和27.0 $\mu\text{g/mL}$, 且有抗污损作用、 EC_{50} 为7.7 $\mu\text{g/mL}$ ^[42]。 *A. sydowii* PSU-F154 代谢产生倍半萜 aspergillusenes A (**108**)、B (**109**)和




图 2. 化合物93–173的结构

Figure 2. Structures of compounds 93–173.

methyldsydonic acid (**110**), 以及氧杂蒽酮 aspergillusones A(**111**)和B(**112**)^[43]。 *A. versicolor* LCJ-5-4代谢产生环五肽versicoloritides A–C (**113–115**)、地衣酚四聚体tetraorcinol A(**116**)、内酯versicolactones A、B(**117**、**118**)^[44]以及喹唑啉酮生物碱cottoquinazolines B–D(**119–121**)^[45], 其中化合物**116**有弱的DPPH自由基清除活性(IC_{50} 为67 $\mu\text{mol/L}$), 化合物**121**对白色念珠菌有中等抑菌活性(MIC 22.6 $\mu\text{mol/L}$); 化合物**116**^[46]、**121**^[47]被NPR选为热点化合物; 经过全合成研究, 化合物versicolactones A和B的结构分别修正为**122**和**123**^[48]。烟曲霉*A. fumigatus*代谢产生新吡啶生物碱**124**和**125**^[49]。二酮哌嗪生物碱spirotryprostatin F(**126**)^[50]和fumiquinazoline K(**127**)以及萜类化合物**128**^[51]来自另一株烟曲霉*A. fumigatus* KMM 4631, 其中化合物**126**在较低浓度(0.1–1.0 $\mu\text{mol/L}$)时可以促进大豆、荞麦和小麦发芽^[50]。吡啶生物碱cyclotryprostatin E(**129**)来自萨氏曲霉*A. sydowii* SCSIO 00305^[52]。内生曲霉*Aspergillus* sp.代谢产生二倍半萜ophiobolin O(**130**)和6-*epi*-ophiobolin O(**131**), 对P388均有较强的细胞毒活性, IC_{50} 分别为4.7和9.3 $\mu\text{mol/L}$ ^[53], 且化合物**130**可以使MCF-7周期停滞于G₀/G₁期(IC_{50} 为17.86 $\mu\text{mol/L}$)、并通过MAPK途径使MCF-7呈现时间和剂量依赖的细胞凋亡^[54]。土曲霉菌*A. terreus* SCSGAF0162代谢产生asperterrestide A(**132**)、terremide C(**133**)和aspernolide E(**134**)(图2), 化合物**132**对组织细胞淋巴瘤细胞U937和急性淋巴母细胞白血病细胞MOLT-4的 IC_{50} 值分别是6.4和6.2 $\mu\text{mol/L}$, 对流感病毒H1N1和H3N2的 IC_{50} 值分别是15.0 $\mu\text{mol/L}$ 和8.1 $\mu\text{mol/L}$ ^[55]。

曲霉菌*Aspergillus* sp. XS-20090066代谢产生吡啶生物碱17-*epi*-notoamides Q、M(**135**、**136**)和苯醚衍生物cordyols D、E(**137**、**138**)^[56]。曲霉菌*A. elegans* ZJ-2008010代谢产生4'-OMe-asperphenamate (**139**)和细胞松弛素aspochalasins A₁和Z₂₄(**140**、

141), 其中化合物**139**对表皮葡萄球菌*S. epidermidis*表现出了选择性的抑制活性(MIC值为10 $\mu\text{mol/L}$)^[57]。曲霉菌*Aspergillus* sp. SCSGAF 0076代谢产生了大环内酯aspergillide D(**142**)^[58]。赤散囊菌*Eurotium rubrum* SH-823代谢产生硫代物eurothiocins A和B(**143**和**144**), 抑制 α -糖苷酶的 IC_{50} 值分别为17.1 $\mu\text{mol/L}$ 和42.6 $\mu\text{mol/L}$ ^[59]。杂色曲霉*A. versicolor*代谢产生核苷类化合物**145**和**146**, 对表皮葡萄球菌有选择性抑制作用、MIC值为12.5 $\mu\text{mol/L}$, 对卤虫有致死活性, LC_{50} 值为8.4 $\mu\text{g/mL}$ ^[60]。曲霉菌*A. flavipes*代谢产生脑苷脂flavicerebrosides A和B(**147**和**148**)(图2), 均对KB细胞有细胞毒活性, IC_{50} 值分别为20.7 $\mu\text{g/mL}$ 和14.3 $\mu\text{g/mL}$ ^[61]。

1.3 其它动物来源的曲霉天然产物

Fumiquinazolines A–C(**149–151**)来自烟曲霉*A. fumigatus* OUPS-T106B-5, 有中等细胞毒活性^[12](Numata et al. 1992), 其结构通过全合成确证^[62–63]; 该菌株还产生fumiquinazolines D–G(**152–155**)^[64]、cephalimysin A(**156**)^[65]及cephalimysins B–D(**157–159**)^[66], 其中化合物**149**、**150**及**152–155**对P388细胞有细胞毒活性, ED_{50} 分别为6.1、16.0、13.5、13.8、14.6和17.7 $\mu\text{g/mL}$ ^[64], 化合物**156**对P388和HL-60细胞的 IC_{50} 值分别为15.0 nmol/L和9.5 nmol/L^[65], 化合物**158**和**159**对HL-60的 IC_{50} 分别为58.4和48.7 $\mu\text{mol/L}$ ^[66]。曲霉*A. terreus* OUCMDZ-1925代谢产生rubrolides R(**160**)和S(**161**), 抑制K562细胞的 IC_{50} 分别为12.8和10.9 $\mu\text{mol/L}$, 化合物**161**还具有抗H1N1病毒的活性、 IC_{50} 为87.1 $\mu\text{mol/L}$, 化合物**160**具有ABTS和DPPH自由基清除活性, IC_{50} 分别为1.33 mmol/L和43.4 $\mu\text{mol/L}$ ^[67]。曲霉*Aspergillus* sp. MF275代谢产生himeic acids A–C(**162–164**)^[68]和himeic acids E–G(**165–167**)^[69](图2), 其中化合物**162**具有泛素激活酶E1抑制活性, 50 $\mu\text{mol/L}$ 时的抑制率为65%^[68]。

曲霉*Aspergillus* sp. MF297-2代谢产生notoamides A–D(**168–171**)(图2), 化合物**168–170**对Hela及

L1210细胞的 IC_{50} 值为22–52 $\mu\text{g/mL}$ ^[70]；该菌株还代谢产生notoamides F–K(**172–177**)(图3)，其中**175**对Hela有弱活性、 IC_{50} 为21 $\mu\text{g/mL}$ ^[71]。随后，又从该菌株的代谢产物中相继分离鉴定了notoamide E(**178**)和notoamides E₂–E₄(**179–181**)^[72]、(-)-versicolamide B(**182**)和notoamides L–N(**183–185**)^[73]、notoamides O–R(**186–189**)^[74]以及notoamide S(**190**)^[75]。具有倍半萜母核的吡啶生物碱pileotin A(**191**)则来自烟曲霉*A. fumigatus* OUPS-N138^[76]。变色曲霉*A.*

versicolor OUPS-N136代谢产生吡啶二萜anthcolorins A–F(**192–197**)(图3)，其中**193–195**对P388细胞的 IC_{50} 值在2.2–8.5 $\mu\text{mol/L}$ 之间^[77]。

Spiculisporic acids B–D(**198–200**)来自内生曲霉*Aspergillus* sp. HDf2^[78]。环肽clavatustides A、B(**201、202**)^[79]和C(**203**)^[80]来自1株棒曲霉*A. clavatus* C2WU，其中**201**和**202**呈现剂量依赖的肝癌细胞系(HCC)增殖抑制活性。此外，**201**、**202**还可将人肝癌细胞HepG-2的细胞周期阻滞在




图 3. 化合物174–251的结构

Figure 3. Structures of compounds 174–251.

G₁期^[79]。化合物**204**、spirotryprostatins C-E (**205–207**)、fumitremorgin B的衍生物(**208**、**209**)和13-oxoverruculogen(**210**)来自一株烟曲霉 *A. fumigatus*, 均有细胞毒活性, 尤其是化合物**207**对MOLT-4、HL-60和A549细胞的IC₅₀分别为3.1、2.3、3.1 μmol/L, **208**对HL-60、BEL-7402细胞的IC₅₀分别为3.4、7.0 μmol/L, **209**和**210**对HL-60细胞的IC₅₀分别为5.4 μmol/L和1.9 μmol/L^[81]。另一株烟曲霉 *A. fumigatus* WFZ-25代谢产生螺内酰胺 pseurotins A₁和A₂(**211**和**212**)^[82]。Yanuthones A-E(**A213–A217**)、1-hydroxyyanuthones A、C(**218**、**219**)及22-deacetylyanuthone A(**220**)(图3)产自黑曲霉 *A. niger* F97S11^[83]。

曲霉菌 *Aspergillus* sp. MF297代谢产生多聚乙酰aspermytin A(**221**), 在50 μmol/L时**221**可诱导小鼠嗜铬细胞瘤PC-12细胞神经轴突生长^[84]。黄曲霉 *A. flavus* OUCMDZ-2205代谢产生吡啶二萜**222**、**223**和异香豆素**224**, 其中化合物**222**对金黄色葡萄菌的MIC值为20.5 μmol/L, 在10 μmol/L时**222**和**223**可将A549细胞阻滞在S期, 化合物**222**对PKC-β的IC₅₀值为15.6 μmol/L^[85]。柄曲霉 *A. flavipes* Z-4代谢产生环肽**225**^[86](图3)。

2 植物来源的曲霉天然产物

2.1 海藻来源的曲霉天然产物

Tetrahydrobostrycin(**226**)和1-deoxytetrahydrobostrycin(**227**)来自曲霉 *Aspergillus* sp. 05F16^[87]。倍半萜 insulicolide A(**228**)来自胰岛曲霉 *A. insulicola*^[88]。杂色曲霉 *A. versicolor* CNC 327代谢产生化合物**229–232**(图3), 其中**229**表现出较强细胞毒活性, 对HCC-2998、HCT-116、BT-549和SNB-75细胞的LC₅₀分别为0.53、0.44、0.27和0.44 μg/mL, 其对5株肾肿瘤细胞(786-0、ACHN、CAK-1、TK-10和UO-31)表现出选择性抑制活性、LC₅₀约为0.47–0.57 μg/mL^[89]。曲霉 *Aspergillus* sp.代谢产生

mactanamide(**233**), 具有抑制白色念珠菌活性^[90]。Parasitenone(**234**)来自一株曲霉 *A. parasiticus* MFA153^[91]。手性双吡咯并萜terreusinone(**235**)产自土曲霉 *A. terreus* MFA 460, 具有防护紫外线A的活性, ED₅₀值为70 μmol/L^[92]。曲霉菌 *Aspergillus* sp. MFA 212代谢产生二酮哌嗪golmaenone(**236**), 具有防护紫外线A的活性(ED₅₀值为90 μmol/L)和DPPH自由基清除活性, IC₅₀值为20 μmol/L^[93]。Sydowins A、B(**237**、**238**)来自萨氏曲霉 *A. sydowii*^[94]。从曲霉 *Aspergillus* sp. MFB024的代谢产物中发现1个多氧萜氢衍生物**239**(chlorofusarielin B), 其对金黄色葡萄球菌、甲氧西林耐药葡萄球菌、多药耐药葡萄球菌具有一定的抑制作用, 其MIC均为62.5 μg/mL^[95]。Nigerasperones A–C(**240–242**)为黑曲霉 *A. niger* EN-13的产物, 化合物**242**对白色念珠菌的抑菌圈为9 mm(两性霉素的抑菌圈为12 mm)、对DPPH自由基的清除率为41.6%^[96]。此菌还代谢产生asperamides A和B(**243**和**244**)^[97]、ergosterimide(**245**)^[98]、**246**^[99]、isopyrophen(**247**)和aspergillusol(**248**)^[100], 其中化合物**243**对白色念珠菌的抑菌圈为12 mm^[97]。

化合物**249**、**250**^[101]和iso-α-CPA(**251**)(图3)^[102]产自黄曲霉 *A. flavus* sc-f-3, 化合物**251**对A549细胞的IC₅₀值为42.2 μmol/L^[102]。赭曲霉 *A. ochraceus* EN-31代谢产生2-hydroxycircumdatin C(**252**)^[103]、7-nor-ergosterolide(**253**)和化合物**254**、**255**^[104](图4), 化合物**252**具有明显的DPPH自由基清除作用, 其IC₅₀值为9.9 μmol/L^[103]。吡啶二萜 asporyzin A–C(**256–258**)^[105]和甾类 asporyergosterol(**259**)^[106]来自米曲霉 *A. oryzae* cf-2, 其中化合物**A258**有较强的大肠杆菌抑制作用(每孔30 μg给药时, 抑菌圈为8 mm)^[105]。JBIR-81(**260**)和JBIR-82(**261**)来自曲霉 *Aspergillus* sp. SpD081030G1f1, 是有效的自由基清除剂(对N18-RE-105细胞的L-谷氨酸毒性的EC₅₀分别为0.7和1.5 μmol/L, 强于对照组的8.8 μmol/L)^[107]。脑苷脂类 flavusides




图 4. 化合物252–331的结构

Figure 4. Structures of compounds 252–331.

A(262)和B(263)来自黄曲霉*A. flavus*, 抑制金黄色葡萄球菌的MIC为15.6 $\mu\text{g/mL}$, 对MRSA的MIC约为31.2 $\mu\text{g/mL}$ ^[108]。脂肪酸264和甾体265来自另一株黄曲霉*A. flavus* cf-5^[109]。肉色曲霉*A. carneus* KMM4638代谢产生吡啶生物碱carneamides A–C(266–268)、喹唑酮生物碱carnequinazoles A–C(269–271)、芳基糖苷carnemycins A、B(272、273)及倍半萜274^[110]。喹唑啉酮衍生物275、276和

二苯醚衍生物277也来自该菌^[111]。二萜asperolides A–C(278–280)^[112]、wentiquinone C(281)和282^[113](图4)来自温特曲霉*A. wentii* EN-48, 其中化合物278和279对多种肿瘤细胞有弱的细胞毒活性(IC_{50} 为35–97 $\mu\text{mol/L}$)、化合物282有明显的DPPH自由基清除活性(IC_{50} 为5.2 $\mu\text{g/mL}$)。

化合物283来自一株变色曲霉*A. versicolor*, 对枯草芽孢杆菌、蜡状芽孢杆菌和金黄色葡萄球

菌有中等强度的抑菌活性(每孔100 μg 给药时的抑菌圈分别为11、12和14 mm)^[114]。萜类化合物1,2-dihydroterretonin F(**284**)、(6 α)-21-deoxyophiobolin G(**285**)、(6 α)-16,17-dihydro-21-deoxyophiobolin G(**286**)、ophiobolin U(**287**)、ophiobolin V(**288**)、ophiobolin W(**289**)和**290**、**291**(图4)来自焦曲霉*A. ustus* cf42, 其中化合物**287**对大肠杆菌和金黄色葡萄球菌均有抑菌活性、每孔30 μg 给药时的抑菌圈分别为15和10 mm^[115]。Asperversin A(**292**)和**293**产自变色曲霉*A. versicolor* pt20^[116]。花斑曲霉*A. versicolor* dl29代谢产生倍半萜albican-11,14-diol(**294**)^[117]和生物碱aspeverin(**295**)^[118], 其中化合物**294**对卤虫的 LC_{50} 为35.0 $\mu\text{g}/\text{mL}$, 且为大肠杆菌和金黄色葡萄球菌的抑制剂、30 $\mu\text{g}/\text{孔}$ 时的抑菌圈分别为7和10.3 mm^[117], 而化合物**295**对赤潮异弯藻*Heterosigma akashiwo*的 EC_{50} 值分别为6.3 $\mu\text{g}/\text{mL}$ (24h)和3.4 $\mu\text{g}/\text{mL}$ (96h)^[118]。蒽醌6,8-di-*O*-methylaverantin(**296**)来自变色曲霉*A. versicolor* EN-7^[119]。蜡叶散囊菌*E. herbariorum* HT-2代谢产生了吡啶二酮哌嗪二聚体cristatumin E(**297**), 其对K562细胞的 IC_{50} 值为8.3 $\mu\text{mol}/\text{L}$, 对产气杆菌*Enterobacter aerogenes*和大肠杆菌*Escherichia coli*的MIC值分别为44.0和44.0 $\mu\text{mol}/\text{L}$ ^[120]。赭曲霉*A. ochraceus* Jcma1F17代谢产生倍半萜6 β ,9 α -dihydroxy-14-*p*-nitrobenzoylcinnamolide(**298**), 对10种人体肿瘤细胞(H1975、U937、K562、BGC-823、Molt-4、MCF-7、A549、Hela、HL60和Huh-7)的 IC_{50} 为1.95–6.12 $\mu\text{mol}/\text{L}$, 对H3N2病毒和EV71病毒的 IC_{50} 分别为17.0和9.4 $\mu\text{mol}/\text{L}$ ^[121]。2株曲霉菌*Aspergillus* sp. BM-05和BM-05ML共培养产生环三肽psychrophilin E(**299**)(图4), 对HCT-116细胞的 IC_{50} 值为28.5 $\mu\text{mol}/\text{L}$ ^[122]。曲霉菌*A. pseudodeflectus*代谢产生pseudodeflectusin(**300**), 对NUGC-3、HeLA-S3和HL-60均有细胞毒活性, 其中对HL-60细胞的 IC_{50} 值为39 $\mu\text{mol}/\text{L}$ ^[123]。

1997年, Fenical等从曲霉*Aspergillus* sp. CNC139的代谢产物中分离得到二酮哌嗪类halimide(**301**), 其对HCT116和A2780细胞有较强的细胞毒活性, IC_{50} 分别为1 $\mu\text{mol}/\text{L}$ 和0.8 $\mu\text{mol}/\text{L}$ ^[3-4]。同时Kano等也从焦曲霉*A. ustus* NSC-F038的代谢产物中分离得到, 将其命名为phenylahistin, 并发现(-)-phenylahistin(**302**)才是真正的活性成分, 对P388的 IC_{50} 为0.35 $\mu\text{mol}/\text{L}$, 而且1 $\mu\text{mol}/\text{L}$ 时可以将该细胞阻滞在G₂/M期^[124], 良好的生物活性使其成为先导结构。之后多个课题组通过全合成出^[125-126], 构效关系研究筛选出plinabulin(NPI-2358)(**303**)^[5], 作为肿瘤细胞的血管分裂剂进入了II期临床研究^[127-129]。目前, plinabulin已经结束其II期临床研究, 并于2015年第三季度开始在美国和中国进行其III期临床研究, 用于治疗转移性的晚期非小细胞肺癌^[6]。Gerwick所列的20个海洋药物中(包括7个上市药和13个临床药物), **303**是唯一的海洋曲霉属真菌来源的药物^[7]。近期, Hayashi小组对plinabulin进行结构改造得到活性更好的化合物KPU-300(**304**)(图4), **304**对HT-29细胞的 IC_{50} 为7.0 nmol/L, 可以有效的与微管蛋白结合($K_d = 1.3 \mu\text{mol}/\text{L}$), 诱导微管解聚^[130]。

2.2 红树林来源的曲霉天然产物

黑曲霉*A. niger* LL-LV3020代谢产生pyranonigrin A(**305**)^[131]。黄柄曲霉*A. favipes*代谢产生cytochalasins Z₁₆–Z₂₀(**306**–**310**)^[132]。赤散囊菌*Eurotium rubrum* QEN-0407-G2代谢产生蒽酮衍生物eurorubrin(**311**)和**312**–**314**, 其中化合物**311**显示了中等强度的DPPH自由基清除活性, IC_{50} 值为44.0 $\mu\text{mol}/\text{L}$ ^[133]。苯并- γ -吡喃酮二聚体rubasperones A–C (**315**–**317**)^[134]和rubasperones D–E(**318**–**321**)^[135]分离自塔宾曲霉*A. tubingensis* GX1-5E。Nigerapyrones A–H(**322**–**329**)和已知物asnipyrones A和B来自黑曲霉*A. niger* MA132, 其中已知化合物asnipyrones A和B的结构分别被修正为**330**和

331(图4); 化合物**323**对HepG2细胞的 IC_{50} 为62 $\mu\text{mol/L}$, **326**对SW1990、DMA-MB-231和A549细胞的 IC_{50} 分别为38、48和43 $\mu\text{mol/L}$ ^[136]。单萜 acetoxhydroaustin B(**332**)和1,2-dihydroacetoxhydroaustin B(**333**)(图5)来自 *Aspergillus* sp. 085241B^[137]。黄曲霉 *A. flavus* 092008代谢产生黄曲霉毒素aflatoxin B_{2b}(**334**), 其对大肠杆菌、枯草芽孢杆菌和产气杆菌有中等的

抑菌活性(MIC分别为22.5、1.7和1.1 $\mu\text{mol/L}$), 对A549、K562和L-02细胞的 IC_{50} 分别为8.1、2.0和4.2 $\mu\text{mol/L}$ ^[138]。Aspergillumarins A(**335**)和B(**336**)产自曲霉 *Aspergillus* sp., 在50 $\mu\text{g/mL}$ 时对金黄色葡萄球菌和芽孢杆菌有抑制活性^[139]。

构巢曲霉 *A. nidulans* MA-143代谢产生aniduquinolones A–C(**337–339**)、6-deoxyaflaquinolone E(**340**)、isoaflaquinolone E(**341**)、14-




图 5. 化合物332–423的结构

Figure 5. Structures of compounds 332–423.

hydroxyaflaquinolone F(**342**)和已知物aflaquinolone A(**343**)(图5); 化合物**338**、**339**和**343**具有抗卤虫活性, LD_{50} 值分别是7.1、4.5和5.5 $\mu\text{mol/L}$ ^[140]。喹唑啉酮生物碱aniquinazolines A–D(**344**–**347**)也产自该菌, 对卤虫的 LD_{50} 分别为1.27、2.11、4.95 和 3.42 $\mu\text{mol/L}$ ^[141]。黄柄曲霉*A. flavipes* AIL8代谢产生flavipesins A(**348**)和B(**349**), **348**对金黄色葡萄球菌和枯草芽孢杆菌的MIC分别为8.0 $\mu\text{g/mL}$ 和 0.25 $\mu\text{g/mL}$ ^[142]。二倍半萜asperterpenols A(**350**)和 B(**351**)来自一株内生曲霉*Aspergillus* sp. 085242, 二者对乙酰胆碱酯酶的 IC_{50} 值分别为2.3和3.0 $\mu\text{mol/L}$ ^[143]。曲霉菌*Aspergillus* sp. 16-5c代谢产生二倍半萜asperterpenoid A(**352**)(图5), 它对分枝杆菌*Mycobacterium tuberculosis*蛋白酪氨酸磷酸酶的 IC_{50} 值为2.2 $\mu\text{mol/L}$ ^[144]。土曲霉*A. terreus* GX7-3B代谢产生了噻吩萜醌衍生物**353**^[145]。

2.3 其它植物来源的曲霉天然产物

吡啶单萜speradine A(**354**)产自溜曲霉*A. tamarii* M143, 抑制 Ca^{2+} -ATP酶的 IC_{50} 为8 $\mu\text{mol/L}$ 、抑制组蛋白去乙酰化酶的 IC_{50} 为100 $\mu\text{g/mL}$ 、抑制藤黄微球菌*Mycrococcus luteus*的MIC 16.7 $\mu\text{g/mL}$ ^[146]。6-methoxyspirotryprostatin B(**355**)、18-oxotryprostatin A(**356**)、14-hydroxyterezine D(**357**)、14-norpseurotin A(**358**)和**359**(图5)来自萨氏曲霉*A. sydowi* PFW1-13, 其中化合物**355**–**357**对A549细胞的 IC_{50} 分别为8.29、1.28和7.31 $\mu\text{mol/L}$, 化合物**358**和**359**对大肠杆菌、枯草杆菌和溶壁微球菌的MIC分别为3.74、14.97、7.49 $\mu\text{mol/L}$ 和 10.65、5.33、10.65 $\mu\text{mol/L}$ ^[147]。

3 海泥、海水等来源的曲霉天然产物

3.1 海泥来源的曲霉天然产物

肉色曲霉*A. carneus* MST-MF156代谢产生aspergillicins A–E(**360**–**364**)(图5), 对捻转血矛线

虫*Haemonchus contortus*具有细胞毒性(LD_{99} 为 25–50 $\mu\text{g/mL}$)^[148]。糖苷类化合物**365**产自变异曲霉*A. varians* KMM 4630, 其在10.0 $\mu\text{g/mL}$ 时对海胆胚胎有毒性^[149]。2,3-dimethoxyosate(**366**)产自曲霉*Aspergillus* sp. B-F-2, 其对K562细胞的 IC_{50} 为 76.5 $\mu\text{mol/L}$, 在100 $\mu\text{mol/L}$ 时可以诱导细胞凋亡、使细胞停滞在S期^[150]。烟曲霉*A. fumigates* 030402d代谢产生11-*O*-methylpseurotin A(**367**), 对Hof1 缺失的酵母菌*Saccharomyces cerevisiae* 的抑菌圈直径为9 mm^[151]。白曲霉*A. candidus* RF-5672代谢产生terprenin(**368**)、3-methoxyterprenin (**369**)和4"-deoxyterpren(**370**), 其抑制CoA-A 介导的小鼠脾淋巴细胞的 IC_{50} 依次为1.2、2.0和5.6 ng/mL, 抑制LPS介导的小鼠脾淋巴细胞的 IC_{50} 分别为4.5、8.0、15.6 ng/mL^[152]。炭黑曲霉*A. carbonarius* WZ-4-11代谢产生carbonarones A (**371**)和B(**372**)^[153]及化合物**373**–**374**^[154], 其中**371**和**372**对K562细胞的 IC_{50} 值分别为56.0和27.8 $\mu\text{g/mL}$ ^[153], **373**和**374**对结核分歧杆菌*Mycobacterium tuberculosis* H37Rv的MIC依次为43.0和25.1 $\mu\text{mol/L}$ ^[154]。烟曲霉*A. fumigates* Fres代谢产生胶霉素**375**^[155]和二酮哌嗪**376**–**378**^[156]。变色曲霉*A. versicolor* MST-MF495代谢产生cottoquinazoline A(**379**)和cotteslosins A(**380**)和B(**381**)^[157]。氧杂螺内酰胺azaspirofurans A(**382**)和B(**383**)来自萨氏曲霉*A. sydowi* D2-6, **382**对A549的 IC_{50} 10 $\mu\text{mol/L}$ ^[158]。二酮哌嗪 azonazine(**384**)来自胰岛曲霉*A. insulicola* 088708a, 有抗炎作用, 其对NF- κ B的 IC_{50} 为8.37 $\mu\text{mol/L}$ ^[159]。杂萜asperdemin(**385**)来自变色曲霉*A. versicolor*, 有溶血作用, EC_{50} 1.15 mmol/L^[160]。环肽unguisin E(**386**)和deoxyapoaranotin(**387**)分别来自曲霉*Aspergillus* sp. AF119^[161]和变色曲霉*A. versicolor* KMD 901^[162], *Aspergillus* sp. AF119还代谢产生barceloneic lactones B(**388**)和C(**389**)和5'-hydroxychlorflavonin(**390**)^[163]以及terphyl acid(**391**)、terphyl diacid(**392**)^[164](图5)。

Protulactones A、B(**393**、**394**)^[165]及protuboxepins A、B(**395**、**396**)、protubonines A、B(**397**、**398**)(图5)来自*Aspergillus* sp. SF-5044^[166], 该菌还代谢产生aflaquinolones A–G(**399–405**)^[167]。*A. versicolor* ZLN-60产生环戊肽versicotides A(**406**)、B(**407**)^[168]和异戊二烯化的二苯醚衍生物diorcinsols B–E(**408–411**)^[169], 其中化合物**410**对Hela和K562细胞的 IC_{50} 值分别为31.5和48.9 $\mu\text{mol/L}$ 、**411**对Hela细胞的 IC_{50} 值为36.5 $\mu\text{mol/L}$ ^[169]。台中曲霉*A. taichungensis* ZHN-7-07代谢产生prenylterphenyllins A–C(**412–414**)、4"-dehydro-3-hydroxyterphenyllin(**415**)及prenylcandidusins A–C(**416–418**), 其中化合物**412**对HL-60和A549细胞的 IC_{50} 分别为1.5和8.3 $\mu\text{mol/L}$ 、**415**和**417**对P388细胞的 IC_{50} 分别为2.7和1.6 $\mu\text{mol/L}$ ^[170]。补身烷倍半萜**419–423**(图5)来自一株焦曲霉*A. ustus*, 化合物**422**抑制P388的 IC_{50} 为8.7 $\mu\text{mol/L}$ ^[171]。Prenylcyclotryprostatin B(**424**)、20-hydroxycyclotryprostatin B(**425**)、9-hydroxyfumitremorgin C(**426**)、6-hydroxytryprostatin B(**427**)和spirogliotoxin (**428**)(图6)来自烟曲霉*A. fumigates* YK-7, 化合物**424**和**426**对U937细胞的 IC_{50} 分别为25.3 $\mu\text{mol/L}$ 和18.2 $\mu\text{mol/L}$ ^[172]。土曲霉*A. terreus* A8-4代谢产生7"-hydroxybutyrolactone III(**429**)和terretriones A–C(**430–432**)^[173]; 三肽presclerototide F(**433**)来自胰岛曲霉*A. insulicola* 088708aZA^[174]; 萜烷衍生物decumbenone C(**434**)来自硫色曲霉*A. sulphureus* KMM 4640, 对人体黑色素瘤SK-MEL-5的细胞毒活性 IC_{50} 为0.9 $\mu\text{mol/L}$ ^[175]。二酮哌嗪brevianamides S–V(**435–438**)来自变色曲霉*A. versicolor* MF030, 化合物**435**具有选择性地抑制结核分支杆菌*Mycobacterium bovis*减毒株BCG的活性(MIC为6.25 $\mu\text{g/mL}$), 可能发展为抗结核杆菌先导药物^[176]。焦曲霉*A. ustus* 094102代谢产生倍半萜ustusols A–C(**439–441**)和ustusolates A–E(**442–446**)以及香豆素ustusoranes A–F(**447–452**), 其中**446**和**451**对HL-60的 IC_{50} 值分别为9.00 $\mu\text{mol/L}$ 和0.13 $\mu\text{mol/L}$, **444**对A549细胞的 IC_{50} 为10.5

$\mu\text{mol/L}$ ^[177]。外消旋的螺环生物碱effusin A(**453**)和dihydrocryptoechinulin D(**454**)(图6)分离自赭曲霉*A. effuses* H1-1^[178–179], 其中**454**对P388和HL-60细胞的 IC_{50} 分别为1.83 $\mu\text{mol/L}$ 和4.80 $\mu\text{mol/L}$ 、在100 $\mu\text{mol/L}$ 时可以选择性的抑制拓扑异构酶I的活性^[178]。

曲霉*A. westerdijkiae* DFFSCS013代谢生物碱circumdatins K(**455**)和L(**456**)、5-chlorosclerotiamide (**457**)、10-*epi*-sclerotiamide(**458**)和aspergilliamide B(**459**)(图6), 其中化合物**457**和**458**对K562细胞的 IC_{50} 值分别是44和53 $\mu\text{mol/L}$ ^[180]。三聚的sydowiols A–C(**460–462**)产自一株萨氏曲霉*A. sydowii* MF357, 化合物**460**和**462**对结核分枝杆菌*M. tuberculosis*蛋白酪氨酸磷酸酶PtpA的 IC_{50} 值分别为14 $\mu\text{g/mL}$ 和24 $\mu\text{g/mL}$ 。此外, 化合物**462**对金黄色葡萄球菌的MIC值为12.5 $\mu\text{g/mL}$ ^[181]。棘孢曲霉*A. aculeatus*代谢产生新苯醌aculeatusquinones A–D(**463–466**), 其中**464**和**466**对HL-60、K562和A549细胞的 IC_{50} 值在5.4–76.1 $\mu\text{mol/L}$ 之间^[182]。杂色曲霉*A. versicolor* HDN08-60代谢产生吡啶二酮哌嗪versicamides A–H(**467–474**); 其中**474**对Hela、HCT-116、HL-60和K562细胞的 IC_{50} 分别为19.4、17.7、8.7和22.4 $\mu\text{mol/L}$, **474**也可抑制多种酪氨酸激酶的活性, 10 $\mu\text{mol/L}$ 对KDR、RET和EGFR激酶的抑制率为23%到35%, 对c-Kit的抑制率为60%^[183]。米曲霉*A. oryzae*代谢产生吡啶生物碱speradines B–E(**475–478**), **475**和**478**对Hela细胞的 IC_{50} 均为0.20 mmol/L^[184]; 吡啶生物碱speradines F–H(**479–481**)和circumdatin G(**482**)分别来自米曲霉*A. oryzae*^[185]和褐黄曲霉*A. ochraceus*^[186]。灰绿曲霉*A. glaucus* HB1-19代谢产生aspergentisyls A、B(**483**、**484**)和aspergiodiquinone(**485**)(图6), **483**和**484**具有DPPH自由基清除活性, IC_{50} 值分别为9.3 $\mu\text{mol/L}$ 和17.6 $\mu\text{mol/L}$ ^[187]。蒽醌类衍生物aspergiolide A(**486**)产自同一株菌, 对A549、HL-60、BEL-7402有细胞毒活性^[188]。




图 6. 化合物424-517的结构

Figure 6. Structures of compounds 424-517.

菌核曲霉 *A. sclerotiorum* PT06-1 在高盐寡营养条件下产生环六肽 sclerotides A、B(487、488)^[189], 在高盐富营养条件下则产生环三肽 sclerotiotides A-K(489-499)^[190] 和 indole-3-ethenamide(500)^[191]; 其中化合物 487 和 488 均显示中等强度的抗白色念

珠菌活性、MIC 值分别为 7.0 $\mu\text{mol/L}$ 和 3.5 $\mu\text{mol/L}$, 化合物 488 对 HL-60 细胞和铜绿假单胞菌有抑制作用、 IC_{50} 和 MIC 值分别为 56.1 和 35.3 $\mu\text{mol/L}$; 化合物 489、490、494 和 497 对白色念珠菌有选择性抑制作用, MIC 分别为 7.5、3.8、

30.0和6.7 $\mu\text{mol/L}$ ，化合物**500**对A549和HL-60细胞的 IC_{50} 分别为3.0和27.0 $\mu\text{mol/L}$ 。土曲霉*A. terreus* PT06-2在盐胁迫条件下代谢产生terremides A、B(**501–502**)和terrelactone A(**503**)，其中**501**对金黄色葡萄球菌的MIC为63.9 $\mu\text{mol/L}$ 、**502**对产气肠杆菌的MIC为33.5 $\mu\text{mol/L}$ ^[192]。烟曲霉*A. fumigates* BM939代谢产生对映异构的tryprostatins A(**504**)和B(**505**)(图6)，浓度分别为50 $\mu\text{g/mL}$ 和12.5 $\mu\text{g/mL}$ 时，两者均能将tsFT-210的细胞周期阻滞于G₂/M期^[193]。

3.2 海水来源的曲霉天然产物

Asperiamide A(**506**)产自曲霉*Aspergillus* sp. MF-34^[194]，asperiamides B(**507**)和C(**508**)则来自黑曲霉*A. niger* MF-16^[195]。曲霉*Aspergillus* sp. MF-93代谢产生asperxanthone(**509**)和asperbiphenyl(**510**)(图6)，均可阻断烟草花叶病毒TMV的复制，0.2 mg/mL浓度下的抑制率分别为62.9%和35.5%^[196]。杂色曲霉*A. versicolor* ZBY-3的新霉素耐药菌株u2n2h3-3代谢产生5-oxo-L-prolinate(**511**)，其对Hela细胞的 IC_{50} 值为49.0 $\mu\text{g/mL}$ ^[197]。柄曲菌素类oxisterigmatocystins A–C(**512–514**)^[198]和二酮哌嗪brevianamide W(**515**)^[199]来自变色曲霉*A. versicolor* CXCTD-06-6a，其中**515**在13.9 $\mu\text{mol/L}$ 时对DPPH的清除率为55%。

3.3 未知来源的曲霉天然产物

二倍半萜aspergilloxide(**516**)和二聚二酮哌嗪**517**(图6)分别来自曲霉*Aspergillus* sp. CNM-713^[200]和黑曲霉*A. niger*^[201]。

4 结论和展望

从1992年Shinggu等首次报道海洋曲霉来源的新天然产物fumiquinazolines A–C^[2](Numata *et al.* 1992)到2014年8月，已发现海洋曲霉来源的新天然产物512个。海洋曲霉天然产物的结构类型多样，包括聚酮、生物碱、萜类、甾体、脂肪酸、肽类及其卤代物和糖苷等；且36%的化合物表现出抗癌(肿瘤细胞毒)、抑菌、抗氧化(自由基清除)和抗寄生虫等生物活性，是发现活性新天然产物和药物先导物的重要资源。

(1) 从海洋曲霉菌的样品来源看，产生新化合物最多的曲霉菌来源或栖息地依次是海泥(149个)、海绵(100个)、其它动物(77个)和海藻(75个)，分别占29%、20%、15%和15%(图7-A)。

(2) 从化合物类型看，化合物最多的类型依次是含氮(265个)、聚酮(153个)和萜甾(96个)，分别占化合物总数的52%、30%和19%(图7-B)。进一步分析其来源，聚酮类化合物产生菌的主要栖息地是海泥和海绵，分别为26%和22%；萜甾化合




图 7. 海洋曲霉天然产物的来源(A)与结构分类(B)

Figure 7. Origin categories (A) and the main structure types (B) of marine-derived *Aspergillus* fungal NPs.

物产生菌的主要栖息地是海绵和海藻, 分别为38%和25%; 含氮化合物产生菌的主要栖息地是海泥, 为35% (图8-A)。进一步结合来源分析, 珊瑚、海藻和海泥来源的曲霉主要代谢产生含氮化合物, 分别为48%、51%和63%; 红树林来源的曲霉主要代谢产生聚酮化合物, 约为64%; 而海绵来源的曲霉天然产物的结构类型相对均衡, 聚酮、含氮化合物及萜甙分别为33%、27%和36% (图8-B)。

(3) 约36%的海洋曲霉天然产物(184个)表现出细胞毒、抗菌、自由基清除和抗寄生虫等多种生物活性(图9-A和表 1), 而肿瘤细胞毒(94个)和抗菌(51个)是其主要的活性, 分别占活性化合物总数的51%和28%。含氮物、聚酮和萜甙是活性化合物的三大类型, 分别占活性化合物总数的57%、29%和20% (图9-B); 卤代物、含氮物、萜甙、聚酮和肽类出现活性化合物的比例分别为各类化合物总数的50%、40%、41%、38%和35%




图 8. 主要结构类型化合物产生菌的栖息地(A)与不同样品来源的曲霉天然产物的结构类群(B)

Figure 8. Habitat categories on the main structures of producing strains (A) and structural categories (B) of NPs from marine-derived *Aspergillus* fungi.




图 9. 海洋曲霉天然产物的活性分类(A)与活性化合物中各结构类型的比例(B)

Figure 9. Bioactive categories of NPs (A) and ratios of the bioactive NPs from structural types (B) of the marine-derived *Aspergillus* fungal origins.

(图10-A); 而海泥和海绵来源的曲霉最容易产生肿瘤细胞毒活性的化合物, 其概率分别为21%和19% (图10-B)。

(4) 我国、其它亚洲国家和欧美国家在海洋天然产物的发展上占据了重要地位, 分别贡献287、108和103个新化合物, 尤其是我国海洋天然产物化学家贡献了57%的海洋曲霉来源的新天然产物 (图11-A)。本文共引用了95篇(占引用文献总数的53%)我国海洋天然产物化学家发表的文章, 是主要的贡献者, 标志我国海洋天然产物的研究具有一定的国际影响力。但其发表在有机化学类或天然产物化学类的主流杂志如*Org. Lett.*、*J. Org.*

Chem.、*Tetrahedron*和*J. Nat. Prod.*分别仅有6、1、2和11篇, 仅占其文章的6.3%、1.1%、2.1%和11.6%, 且未见其在*J. Am. Chem. Soc.*和*Angew. Chem., Int. Ed.*等化学综合类高水平杂志上发表文章(图11-B), 也未见有药进入临床研究。由此可见, 我国曲霉菌海洋天然产物的研究, 应该注重质和国家需求, 而非单纯的量或发表文章。

综上, 海洋曲霉可以产生结构新颖, 活性多样的天然产物, 具有极大的开发潜能, 但是到目前为止仅有1个来源于海洋曲霉的药物, 如何最大限度利用海洋曲霉这类珍贵的微生物资源来开发药物, 是值得科研工作者思考和解决的问题。一




图 10. 不同结构类型化合物的活性率(A)与产生菌来源化合物的活性率(B)

Figure 10. Ratios of active NPs from the structural types (A) and the sources of the *Aspergillus* fungi (B).




图 11. 海洋曲霉天然产物发现者国别(A)和中国学者发表论文的期刊分类(B)

Figure 11. Country categories of the discoverers (A) and publishing Journal categories of the Chinese scholars (B) on the marine-derived fungal NPs

表1. 海洋曲霉来源天然产物(1992–2014)
Table 1. Marine natural products from *Aspergillus* fungi (1992–2014)

Compounds	Producing Strains	Environments source	Bioactivity	References
1, 2	<i>A. cf. ochraceus</i> 941026	<i>Jaspis</i> of <i>Coriacea</i> , Indian-Pacific Ocean	/ ^a	8
3	<i>A. niger</i>	<i>Hyrtios</i> sp., Florida, America	Cytotoxicity	9
4	<i>A. niger</i> 94–1212	<i>Hyrtios proteus</i> , Florida, America	/	10
5–14	<i>A. versicolor</i> (Vuill)Triab	<i>Xestospongia exigua</i> , Bali Island, Indonesia	/	11–12
15–25	<i>A. ostianus</i> TUF 01F313	Unidentified sponge, Pohnpei, Micronesia	15–17: Antibacterial activity, 22–24: Cytotoxicity	13–16
26–28	<i>A. ostianus</i> IBT 12704	Unidentified sponge, Pohnpei, Micronesia	/	17
29–32	<i>Aspergillus</i> sp. CNK-371	Unidentified sponge, Hawaii State	29–31: Cytotoxicity	18
33, 34	<i>A. aculeatus</i> CRI323-04	<i>Xestospongia testudinaria</i> , Phi Phi Island, Thailand	33: α -glucosidase inhibition, Antibacterial activity	19–20
35–37	<i>A. aculeatus</i> CRI322-03	Unidentified sponge, Phi Phi Island, Thailand	/	21
38, 39	<i>A. insuetus</i>	<i>Petrosia ficiformis</i> , Santa Ana Alhambra Nestorius, Spain	Inhibitor of the mammalian mitochondrial respiratory chain	22
40	<i>A. sclerotiorum</i> Huber SP080903f04	<i>Mycale</i> sp. Okinawa Island, Japan	/	23
41–54	<i>A. ustus</i> 8009	<i>Suberites domuncula</i> , The Adriatic Sea	44, 45: Cytotoxicity	24–25
55	<i>A. versicolor</i>	<i>Petrosia</i> sp., Jeju Island, Korea	Cytotoxicity	26
56, 57	<i>Aspergillus</i> sp. fs14	Unidentified sponge, Okinawa Island, Japan	/	27
58–61	<i>A. insuetus</i> OY-207	<i>Psammocinia</i> sp., Israel	58: Antibacterial activity 60: Cytotoxicity	28
62	<i>A. versicolor</i> PF10M	<i>Petrosia</i> sp., Jeju Island, Korea	Cytotoxicity	29
63–69	<i>Aspergillus</i> sp.	<i>Xestospongia testudinaria</i> , South China Sea, China	63–66: Antibacterial activity 67, 69: Cytotoxicity	30–31
70–74	<i>A. unguis</i> CRI282-03	Unidentified sponge CRI282, Thailand	70–72: Aromatase inhibition 70, 71: XXO scavenging activity	32
75	<i>Eurotium cristatum</i> KUFC 7356	<i>Mycale</i> sp. State Beach, Thailand	/	33
76–89	<i>Aspergillus</i> sp.	<i>Tethya aurantium</i> , Mediterranean, Italy	81: Cytotoxicity 88, 89: Antibacterial activity	34–36
90–92	<i>A. versicolor</i> MF359	<i>Hymeniacidon perleve</i> , Bohai, China	92: Antibacterial activity	37
93–99	<i>A. niger</i>	<i>Axinella damicornis</i> , Elba, Italy	93–95: Cytotoxicity	38
100	<i>Aspergillus niger</i> FT–0554	Unidentified sponge, Palau	Inhibit <i>Ascarissuum</i>	39
101, 102	<i>A. terreus</i> HKI0499	<i>Sinularia kavarattiensis</i> , Amanda Pam, India	/	40
103–107	<i>Aspergillus</i> sp.	<i>Dichotella gemmacea</i> , Weizhou Island, China	103: Antibacterial activity 106: Cytotoxicity, fouling resistance	41–42
108–112	<i>A. sydowii</i> PSU-F154	<i>Annella</i> sp., Surat Thani, Thailand	/	43
113–121	<i>A. versicolor</i> LCJ-5-4	<i>Cladiella</i> sp., South China Sea, China	116: DPPH radical scavenging activity 121: Antibacterial activity	44–45
124, 125	<i>A. fumigates</i>	<i>Zoanthus</i> sp., Kagoshima, Japan	/	49

(待续)

(续表1-1)

126–128	<i>A. fumigatus</i> KMM 4631	<i>Sinularia</i> sp., Ostrov Kunashir Island	126: Planta growth Promotion	50–51
129	<i>A. sydowii</i> SCSIO 00305	<i>Verrucella umbraculum</i> , Sanya, China	/	52
130, 131	<i>Aspergillus</i> sp.	<i>Zoanthus</i> sp., Kagoshima, Japan	Cytotoxicity	53–54
132–134	<i>A. terreus</i> SCSGAF0162	<i>Echinogorgia aurantiaca</i> , Sanya, China	132: Cytotoxicity, Antivirus	55
135–138	<i>Aspergillus</i> sp. XS-20090066	<i>Dichotella gemmacea</i> , Xisha Islands, South China Sea	/	56
139–141	<i>A. elegans</i> ZJ-2008010	<i>Sarcophyton</i> sp., Weizhou Island, China	139: Antibacterial activity	57
142	<i>Aspergillus</i> sp. SCSGAF 0076	<i>Melitodes squamata</i> , Sanya, China	/	58
143, 144	<i>Eurotium rubrum</i> SH-823	<i>Sarcophyton</i> sp., South China Sea, China	Anti- α -glycosides	59
145, 146	<i>A. versicolor</i>	<i>Dichotella gemmacea</i> , South China Sea, China	Antibacterial activity, Anti-brine shrimp activity	60
147, 148	<i>A. flavipes</i>	<i>Anthopleura xanthogrammica</i> , Qingdao, China	Cytotoxicity	61
149–156	<i>A. fumigatus</i> OUPS-T106B-5	<i>Pseudolabrus japonicus</i> , Tanabe Bay, Japan	Cytotoxicity	2, 64–65
157–159	<i>A. fumigatus</i> OUPS-T106B-5	<i>Pseudolabrus japonicus</i> , Tanabe Bay, Japan	158, 159: Cytotoxicity	66
160–161	<i>A. terreus</i> OUCMDZ-1925	<i>Chelon haematocheilus</i> , Yellow River estuary, China	DPPH radical scavenging; Cytotoxicity; Anti-virus	67
162–167	<i>Aspergillus</i> sp. MF275	<i>Mytilus edulis</i> , Toyama Bay, Japan	162: Ubiquitin-activating enzyme (E1) inhibitor	68–69
168–190	<i>Aspergillus</i> sp. MF297-2	<i>Mytilus edulis</i> , Japan	170, 175: Cytotoxicity	70–75
191	<i>A. fumigatus</i> OUPS-N138	<i>Toxopneustes pileolus</i> , Japan	/	76
192–197	<i>A. versicolor</i> OUPS-N136	<i>Anthocidaris crassispana</i> , Tanabe Bay, Wakayama, Japan	193–195: Cytotoxicity	77
198–200	<i>Aspergillus</i> sp. HDf2	<i>Anthocidaris crassispina</i> , Hainan, China	/	78
201–203	<i>A. clavatus</i> C2WU	<i>Xenograpsus testudinatus</i> , Taiwan, China	201–202: Cytotoxicity	79–80
204–210	<i>A. fumigatus</i>	<i>S. japonicus</i> , Lingshan Island, Qingdao, China	207–210: Cytotoxicity	81
211–212	<i>A. fumigatus</i> WFZ-25	<i>S. japonicus</i> , Jiaozhou Bay, China	/	82
213–220	<i>A. niger</i> F97S11	<i>Aplidium</i> sp., Fiji	/	83
221	<i>Aspergillus</i> sp. MF297	<i>Mytilus edulis</i> , Toyama Bay, Japan	Cytotoxicity	84
222–224	<i>A. flavus</i> OUCMDZ-2205	<i>Penaeus vannamei</i> , Lianyungang sea area, China	222: Antibacterial activity 222–223: Cytotoxicity	85
225	<i>A. flavipes</i> Z-4	<i>Ligia oceanica</i> , Oceania	/	86
226–227	<i>Aspergillus</i> sp. 05F16	Unidentified marine alga, Indonesia	/	87
228	<i>A. insulicola</i>	Unidentified marine alga, Bahamas	/	88
229–232	<i>A. versicolor</i> CNC 327	<i>Penicillus capitatus</i> , Caribbean	229: Cytotoxicity	89
233	<i>Aspergillus</i> sp.	<i>Sargassum</i> sp., Philippines	233: Antibacterial activity	90
234	<i>A. parasiticus</i> MFA153	<i>Carpopeltis cornea</i> , Korea	/	91
235	<i>A. terreus</i> MFA 460	<i>Halymenia acuminata</i> , Korea	UV-A absorbing activity	92
236	<i>Aspergillus</i> sp. MFA 212	<i>Lomentaria catenata</i> , Ulsan, Korea	UV-A absorbing activity; DPPH radical scavenging	93
237–238	<i>A. sydowii</i>	<i>Acanthophora spicifera</i> , Bay of Bengal India	/	94

(待续)

(续表1-2)

239	<i>Aspergillus</i> sp. MFB024	<i>Sargassum horneri</i> , Korea	Antibacterial activity	95
240–248	<i>A. niger</i> EN-13	<i>Colpomenia sinuosa</i> , Qingdao, China	242: Antibacterial activity; DPPH radical scavenging 243: Antibacterial activity	96–100
249–251	<i>A. flavus</i> c-f-3	<i>Enteromorpha tubulosa</i> , Putian, China	251: Cytotoxicity	101–102
252–255	<i>A. ochraceus</i> EN-31	<i>Sargassum kjellmanianum</i> , Daliancoastline, China	252: DPPH radical scavenging	103–104
256–259	<i>A. oryzae</i> cf-2	<i>Heterosiphonia japonica</i> , Yantai, China	258: Antibacterial activity	105–106
260–261	<i>Aspergillus</i> sp. SpD081030G1f1	<i>Sargassum</i> sp., Ishigaki Island, Japan	DPPH radical scavenging	107
262–263	<i>A. flavus</i>	<i>Codium fragile</i> , Yeosu, Korea	Antibacterial activity	108
264–265	<i>A. flavus</i> cf-5	<i>Corallina officinalis</i> , Yantai, China	/	109
266–277	<i>A. carneus</i> KMM4638	<i>Laminaria sachalinensis</i> , Kunachir Island	/	110–111
278–282	<i>A. wentii</i> EN-48	<i>Sargassum</i> sp., Unknown place	278–282: Cytotoxicity 282: DPPH radical scavenging	112–113
283	<i>A. versicolor</i>	<i>Halimeda opuntia</i> , Egyptian Red Sea (5–8 m)	Antibacterial activity	114
284–291	<i>A. ustus</i> cf42	<i>Codium fragile</i> , Zhoushan Island, China	287: Antibacterial activity	115
292–293	<i>A. versicolor</i> pt20	<i>Sargassum thunbergii</i> , Pingtan Island, China	/	116
294–295	<i>A. versicolor</i> dl29	<i>Codium fragile</i> , Dalian, China	294: Anti-brine shrimp activity, Antibacterial activity 295: Inhibition of <i>H. akashiwo</i>	117–118
296	<i>A. versicolor</i> EN-7	<i>Sargassum thunbergii</i> , Qingdao, China	/	119
297	<i>E. herbariorum</i> HT-2	<i>Enteromorpha prolifera</i> , Qingdao, China	Cytotoxicity, Antibacterial activity	120
298	<i>A. ochraceus</i> Jcma1F17	<i>Coelarthrum</i> sp., Paracel Islands, China	Cytotoxicity, Anti- H3N2 and EV71 activity	121
299	<i>Aspergillus</i> sp. BM-05 and BM-05ML	<i>Sargassum</i> sp., Helgoland, North Sea, Germany	Cytotoxicity	122
300	<i>A. pseudodeflectus</i>	<i>Sargassum fusiform</i> , Miura Peninsula, Japan	Cytotoxicity	123
301	<i>Aspergillus</i> sp. CNC139 <i>A. ustus</i> NSC-F038	<i>Halimeda copiosa</i> , Philippines	Cytotoxicity	3, 4
305	<i>A. niger</i> LL-LV3020	Mangrove wood, Hong Kong, China	/	131
306–310	<i>A. favipes</i>	Mangrove Plant <i>Acanthus ilicifolius</i> , Dongzhai Gang, China		132
311–314	<i>Eurotium rubrum</i> QEN-0407-G2	Marine mangrove plant <i>Hibiscus tiliaceus</i> , Hainan Island, China	311: DPPH radical scavenging	133
315–321	<i>A. tubingensis</i> GX1-5E	Radix of <i>Pongamia pinnata</i> , South China Sea, Guangxi	/	134–135
322–331	<i>A. niger</i> MA132	Mangrove, Hainan, China	323, 326: Cytotoxicity	136
332–333	<i>Aspergillus</i> sp. 085241B	Mangrove, Shankou, Guangxi, China	/	137
334	<i>A. flavus</i> 092008	Mangrove plant, Hainan, China	Cytotoxicity, Antibacterial activity	138
335–336	<i>Aspergillus</i> sp.	<i>Bruguiera gymnorrhiza</i> , South China Sea, China	/	139
337–347	<i>A. nidulans</i> MA-143	<i>Rhizophora stylosa</i> , Unknown place	338–339, 343, 344–347: Anti- brine shrimp activity	140–141
348–349	<i>A. flavipes</i> AIL8	<i>Acanthus ilicifolius</i> , Daya Bay, Shenzhen, China	Antibacterial activity	142
350–351	<i>Aspergillus</i> sp. 085242	Mangrove plant, Guangxi, China	Acetylcholinesterase inhibition	143

(待续)

(续表1-3)

352	<i>Aspergillus</i> sp. 16-5c	Mangrove plant, South China Sea, China	mPTPB inhibition	144
353	<i>A. terreus</i> GX7-3B	<i>Bruguiera gymnoihiza</i> , Guangxi, China	/	145
354	<i>A. tamarii</i> M143	Driftwood, Okinawa Island, Japan	Ca ²⁺ -ATPase inhibition, Antibacterial activity	146
355–359	<i>A. sydowi</i> PFW1-13	Driftwood, Baishamen, Hainan, China	Cytotoxicity, Antibacterial activity	147
360–364	<i>A. carneus</i> MST-MF156	Sediment, Jordan River Bridge, Tasmania, Australia	Antiparasitic activity	148
365	<i>A. varians</i> KMM 4630	Sediment, Sakhalin Island	Cytotoxicity	149
366	<i>Aspergillus</i> sp. B-F-2	Sediment, Behai Bay, China	Cytotoxicity	150
367	<i>A. fumigates</i> 030402d	Sediment (>30 m), Vanuatu	Antimicrobial activity	151
368–370	<i>A. candidus</i> RF-5672	Sediment, Shodo Island, Kagawa Prefecture, Japan	Cytotoxicity	152
371–374	<i>A. carbonarius</i> WZ-4-11	Sediment, Weizhou Island, China	371–372 : Cytotoxicity; 373–374 : Antimicrobial activity	153–154
375–378	<i>A. fumigates</i> Fres	Sediment, Jiaozhou Bay, Qingdao, China	/	155–156
379–381	<i>A. versicolor</i> MST-MF495	Beach sand sample, Cottesloe, Western Australia	/	157
382, 383	<i>A. sydowi</i> D2-6	Sediment, Jiaozhou Bay, Qingdao, China	382 : Cytotoxicity	158
384	<i>A. insulicola</i> 088708a	Sediment, Hawaii	Anti-inflammation	159
385	<i>A. versicolor</i>	Sediment, Sakhalin Bay, Russian	Antihemolysis	160
386	<i>Aspergillus</i> sp. AF119	Sediment, Xiamen beach, China	/	161
387	<i>A. Versicolor</i> KMD 901	Sediment, East Sea, Korea	/	162
388–392	<i>Aspergillus</i> sp. AF119	Sediment, Xiamen beach, China	/	163–164
393–405	<i>Aspergillus</i> sp. SF-5044	Sediment, Dadaepo Beach, Busan, Korea	/	165–167
406–411	<i>A. versicolor</i> ZLN-60	Sediment, Yellow Sea	410, 411 : Cytotoxicity	168–169
412–418	<i>A. taichungensis</i> ZHN-7-07	Root soil of the mangrove plant <i>Acrostichum aureum</i>	412, 415, 417 : Cytotoxicity	170
419–423	<i>A. ustus</i>	Rhizosphere soil of the mangrove <i>Acrostichum aureum</i> , Guangxi, China	422 : Cytotoxicity	171
424–428	<i>A. fumigates</i> YK-7	Sediment, Yingkou, China	424, 426 : Cytotoxicity	172
429–432	<i>A. terreus</i> A8-4	Mangrove-associated marine sediments, Guangxi, China	/	173
433	<i>A. insulicola</i> 088708aZA	Sediment, Hawaii	/	174
434	<i>A. sulphureus</i> KMM 4640	Sediment, Unknown place	Cytotoxicity	175
435–438	<i>A. versicolor</i> MF030	Sediment, Bohai Sea, China	435 : Antitubercular activity	176
439–452	<i>A. ustus</i> 094102	Rhizosphere soil of the mangrove plant <i>Bruguiera gymnorrhiza</i> , Wenchang, Hainan, China	444, 446, 451 : Cytotoxicity	177
453, 454	<i>A. effuses</i> H1-1	Mangrove rhizosphere soil, Fujian, China	454 : Cytotoxicity	178–179
455–459	<i>A. westerdijkiae</i> DFFSCS013	Sediment (–2918 m), South China Sea	457, 458 : Cytotoxicity	180
460–462	<i>A. sydowii</i> MF357	Sediment, Bohai Sea, China	460, 462 : Antimicrobial activity	181

(待续)

(续表1-4)

463–466	<i>A. aculeatus</i>	Sediment, Langqi Island, Fujian, China	464, 466: Cytotoxicity	182
467–474	<i>A. versicolor</i> HDN08-60	Sediment, South China Sea	474: Cytotoxicity; Inhibition of PTKs	183
475–478	<i>A. oryzae</i>	Sediment, Langqi Island, Fujian, China	475, 478: Cytotoxicity	184
479–481	<i>A. oryzae</i>	Sediment, Langqi Island, Fujian, China	/	185
482	<i>A. ochraceus</i>	Sediment, Sea of Japan	/	186
483–485	<i>A. glaucus</i> HB1-19	Mangrove rhizosphere soil, Fujian, China	483, 484: DPPH-radical scavenging	187
486	<i>A. glaucus</i> HB1-19	Mangrove rhizosphere soil, Fujian, China	Cytotoxicity	188
487–500	<i>A. sclerotiorum</i> PT06-1	Putian Sea Salt Field, Fujian, China	487–490, 494, 497: Antimicrobial activity 488, 500: Cytotoxicity	189–191
501–503	<i>A. terreus</i> PT06-2	Putian Sea Salt Field, Fujian, China	501, 502: Antimicrobial activity	192
504, 505	<i>A. fumigates</i> BM939	Sediment (–760 m), Oi River, Japan	Cytotoxicity	193
506	<i>Asperillus</i> sp. MF-34	Sea water, Mei-Zhou Gulf, Fujian, China	/	194
507, 508	<i>A. niger</i> MF-16	Sea water, Quanzhou Gulf, Fujian, China	/	195
509, 510	<i>Aspergillus</i> sp. MF-93	Sea water, Quanzhou Gulf, Fujian, China	Antivirus	196
511	<i>A. versicolor</i> ZBY-3	Sea water (–800 m), Southeast Pacific	Antimicrobial activity; Cytotoxicity	197
512–515	<i>A. versicolor</i> CXCTD-06-6a	Sea water (–800 m), Pacific Ocean	515: DPPH-radical scavenging	198–199
516	<i>Aspergillus</i> sp. CNM-713	Unknown source	/	200
517	<i>A. niger</i>	Unknown source	/	201

^a /: no bioactivity was reported.

方面要通过培养基改造、表观遗传修饰和共培养等手段,继续发现新的活性天然产物;另一方面也是最重要的,是如何开展对现有活性天然产物的成药性研究,而化合物的量依然是制约成药性研究的瓶颈因素。发酵工程、代谢工程是开展微生物活性产物规模化制备、获取足量产物的有效方法,而合成生物学为代谢工程有效而系统的分子生物学工具。由于真菌的基因组较大、酶系复杂,制约着其合成生物学的研究。近年来,曲霉次级代谢产物合成酶系及其相关基因结构、功能的研究取得了很大的进展,多个曲霉的基因组序列被公开。一些曲霉属真菌次级代谢产物的生合成机理被阐明,为曲霉属真菌活性天然产物的规模化制备和工业化生产提供了技术和理论基础。如张立新课题组发现真菌毒素verruculogen中的过

氧桥键是由一个依赖 α -酮戊二酸的单核非血红素酶FtmOx1催化合成的^[202]。降血脂的一线药物洛伐他汀(lovastatin)最初是从曲霉中发现,目前控制洛伐他汀的生物合成酶被发现,并通过曲霉发酵实现工业化生产^[203–204];洛伐他汀的衍生物辛伐他汀(simvastatin)也是一种有效的调血脂药物,经典的合成方法是从洛伐他汀出发,经过水解(得到关键中间体monacolin J)、保护、酰化和脱保护等多步反应得到。Yi Tang等在解析洛伐他汀生物合成中酰化酶的基础上,利用全细胞培养,在大肠杆菌(*Escherichia coli*)过表达酰基转移酶lovD,实现了辛伐他汀的高效合成(转化monacolin J为simvastatin的转化率达到99%以上,且产量达到克级: 4–6 g/L)^[205–206]。这些研究成果为海洋曲霉属真菌药物的开发提供了重要的参考。

致谢

本课题组的王聪博士和马颖娜硕士在本文的写作及文献查阅过程中付出了大量艰辛的劳动, 在此对其贡献表示感谢。

参考文献

- [1] Zhao CY, Zhu TH, Zhu WM. New marine natural products of microbial origin from 2010 to 2013. *Chinese Journal of Organic Chemistry*, 2013, 33(06): 1195-1234 (in Chinese)
赵成英, 朱统汉, 朱伟明. 2010~2013 之海洋微生物新天然产物. *有机化学*, 2013, 33(06): 1195-1234.
- [2] Numata A, Takahashi C, Matsushita T, Miyamoto T, Kawai K, Usami Y, Matsumura E, Inoue M, Ohishi H, Shingu T. Fumiquinazolines, novel metabolites of a fungus isolated from a salt fish. *Tetrahedron Letters*, 1992, 33(12): 1621-1624.
- [3] Fenical W, Jensen PR, Cheng XC. Halimide, a cytotoxic marine natural product, derivatives thereof and therapeutic use in inhibition of proliferation. PCT Int. Appl. WO 9948889. 19990930.
- [4] Fenical W, Jensen P, Cheng X. Halimide, a cytotoxic marine natural product and derivatives thereof. US: US 6069146. 20000530.
- [5] Yamazaki Y, Tanaka K, Nicholson B, Deyanat G, Potts B, Yoshida T, Oda A, Kitagawa T, Orikasa S, Kiso Y. Synthesis and structure- activity relationship study of antimicrotubule agents phenylhistin derivatives with a didehydropiperazine-2, 5-dione structure. *Journal of Medicinal Chemistry*, 2012, 55(3): 1056-1071.
- [6] Beondspring, 2015. <http://www.beyondspringpharma.com/clinical-trials/phase-3/>
- [7] Gerwick WH, Moore BS. Lessons from the past and charting the future of marine natural products drug discovery and chemical biology. *Journal of Biological Chemistry*, 2012, 19(1): 85-98.
- [8] Abrell LM, Borgeson B, Crews P. Chloro polyketides from the cultured fungus (*Aspergillus*) separated from a marine sponge. *Tetrahedron Letters*, 1996, 37(14): 2331-2334.
- [9] Varoglu M, Corbett TH, Valeriote FA, Crews P. Asperazine, a selective cytotoxic alkaloid from a sponge-derived culture of *Aspergillus niger*. *Journal of Organic Chemistry*, 1997, 62(21): 7078-7079.
- [10] Varoglu M, Crews P. Biosynthetically diverse compounds from a saltwater culture of sponge-derived *Aspergillus niger*. *Journal of Natural Products*, 2000, 63(1): 41-43.
- [11] Lin WH, Fu HZ, Li J, Proksch P. Novel chromone derivatives from marine fungus *Aspergillus versicolor* isolated from the sponge *Xestospongia exigua*. *Chinese Chemical Letters*, 2001, 12(3): 235-238.
- [12] Lin WH, Li J, Fu HZ, Proksch P. Four novel hydropyranoindeno-derivatives from marine fungus *Aspergillus versicolor*. *Chinese Chemical Letters*, 2001, 12(5): 435-438.
- [13] Namikoshi M, Negishi R, Nagai H, Dmitrenok A, Kobayashi H. Three new chlorine containing antibiotics from a marine-derived fungus *Aspergillus ostianus* collected in Pohnpei. *The Journal of Antibiotics(Tokyo)*, 2003, 56(9): 755-761.
- [14] Kito K, Ookura R, Yoshida S, Namikoshi M, Ooi T, Kusumi T. Pentaketides relating to aspinonene and dihydroaspyrone from a marine-derived fungus *Aspergillus ostianus*. *Journal of Natural Products*, 2007, 70(12): 2022-2025.
- [15] Kito K, Ookura R, Yoshida S, Namikoshi M, Ooi T, Kusumi T. New cytotoxic 14-membered macrolides from marine-derived fungus *Aspergillus ostianus*. *Organic Letters*, 2008, 10(2): 225-228.
- [16] Kito K, Ookura R, Kusumi T, Namikoshi M, Ooi T. X-ray structures of two stephacidins, heptacyclic alkaloids from the marine-derives fungus *Aspergillus astianus*. *Heterocycles*, 2009, 78(8): 2101-2106.
- [17] Rahbæk L, Breinholt J. Circumdatins D, E, and F: further fungal benzodiazepine analogues from *Aspergillus ochraceus*. *Journal of Natural Products*, 1999, 62(6): 904-905.
- [18] Cueto M, MacMillan JB, Jensen PR, Fenical W. Tropolactones A- D, four meroterpenoids from a marine-derived fungus of the genus *Aspergillus*. *Phytochemistry*, 2006, 67(16): 1826-1831.
- [19] Ingavat N, Dobereiner J, Wiyakrutta S, Mahidol C, Ruchirawat S, Kittakoop P. Aspergillusol A, an α -glucosidase inhibitor from the marine-derived fungus *Aspergillus aculeatus*. *Journal of Natural Products*, 2009, 72(11): 2049-2052.
- [20] Ingavat N, Mahidol C, Ruchirawat S, Kittakoop P. Asperaculin A, a sesquiterpenoid from a marine-derived fungus *Aspergillus aculeatus*. *Journal of Natural Products*, 2011, 74(7): 1650-1652.
- [21] Antia BS, Aree T, Kasetrathat C, Wiyakrutta S, Ekpa OD,

- Ekpe UJ, Mahidol C, Ruchirawat S, Kittakoop P. Itaconic acid derivatives and diketopiperazine from the marine-derived fungus *Aspergillus aculeatus* CRI322-03. *Phytochemistry*, 2011, 72(8): 816-820.
- [22] López-Gresa MP, Cabedo N, Gonzalez-Mas MC, Ciavatta ML, Avila C, Primo J. Terretionins E and F, inhibitors of the mitochondrial respiratory chain from the marine-derived fungus *Aspergillus insuetus*. *Journal of Natural Products*, 2009, 72(7): 1348-1351.
- [23] Motohashi K, Inaba S, Takagi M, Shin-ya K. JBIR-15, a new aspochracin derivative isolated from a sponge-derived fungus, *Aspergillus sclerotiorum* Huber Sp080903f04. *Biosci Biotechnol Biochem*, 2009, 73(8): 1898-1900.
- [24] Liu H, Edrada-Ebel R, Ebel R, Wang Y, Schulz B, Draeger S, Müller WE, Wray V, Lin W, Proksch P. Dimeric sesquiterpenoids from the fungus *Aspergillus ustus* isolated from the marine sponge *Suberites domuncula*. *Journal of Natural Products*, 2009, 72(9): 1585-1588.
- [25] Liu H, RuAngelie RE, Yao BS, Siegfried WEG. Ophiobolin sesterterpenoids and pyrrolidine alkaloids from the sponge-derived fungus *Aspergillus ustus*. *Helvetica Chimica Acta*, 2011, 94(4): 623-631.
- [26] Lee Y, Dang HT, Hong J, Lee CO, Bae K, Kim D, Jung JH. A cytotoxic lipopeptide from the sponge-derived fungus *Aspergillus versicolor*. *Bulletin of the Korean Chemical Society*, 2010, 31(1): 205-208.
- [27] Takagi M, Motohashi K, Shin-ya K. Isolation of 2 new metabolites, JBIR-74 and JBIR-75, from the sponge-derived *Aspergillus* sp. fS14. *The Journal of Antibiotics (Tokyo)*, 2010, 63(7): 393-395.
- [28] Cohen E, Koch L, Thu KM, Rahamim Y, Aluma Y, Ilan M, Yarden O, Carmeli S. Novel terpenoids of the fungus *Aspergillus insuetus* isolated from the Mediterranean sponge *Psammocinia* sp. collected along the coast of Israel. *Bioorganic and Medicinal Chemistry*, 2011, 19(22): 6587-6593.
- [29] Lee Y, Dang HT, Li J, Zhang P, Hong J, Lee CO, Jung JH. A cytotoxic fellutamide from the sponge-derived fungus *Aspergillus versicolor*. *Bulletin of the Korean Chemical Society*, 2011, 32(10): 3817-3820.
- [30] Li D, Xu Y, Shao CL, Yang RY, Zheng CJ, Chen YY, Fu XM, Qian PY, She ZG, de Voogd NJ, Wang CY. Antibacterial bisabolane-type sesquiterpenoids from the sponge-derived fungus *Aspergillus* sp. *Marine Drugs*, 2012, 10(1): 234-241.
- [31] Sun L, Shao C, Chen J, Guo Z, Fu X, Chen M, Chen Y, Li R, de Voogd NJ, She Z. New bisabolane sesquiterpenoids from a marine-derived fungus *Aspergillus* sp. isolated from the sponge *Xestospongia testudinaria*. *Bioorganic and Medicinal Chemistry Letters*, 2012, 22(3): 1326-1329.
- [32] Sureram S, Wiyakrutta S, Ngamrojanavanich N, Mahidol C, Ruchirawat S, Kittakoop P. Depsidones, aromatase inhibitors and radical scavenging agents from the marine-derived fungus *Aspergillus unguis* CRI282-03. *Planta Medica*, 2012, 78(06): 582-588.
- [33] Gomes NM, Dethoup T, Singburauodom N, Gales L, Silva A, Kijjoa A. Eurocristatine, a new diketopiperazine dimer from the marine sponge-associated fungus *Eurotium cristatum*. *Phytochemistry Letters*, 2012, 5(4): 717-720.
- [34] Zhou Y, Má ndi A, Debbab A, Wray V, Schulz B, Müller WEG, Lin W, Proksch P, Kurtá n T, Aly AH. New austalides from the sponge-associated fungus *Aspergillus* sp. *European Journal of Organic Chemistry*, 2011, 2011(30): 6009-6019.
- [35] Zhou Y, Debbab A, Má ndi A, Wray V, Schulz B, Müller WEG, Kassack M, Lin W, Kurtá n T, Proksch P. Alkaloids from the sponge-associated fungus *Aspergillus* sp.. *European Journal of Organic Chemistry*, 2013, 2013(5): 894-906.
- [36] Zhou Y, Debbab A, Wray V, Lin W, Schulz B, Trepos R, Pile C, Hellio C, Proksch P, Aly AH. Marine bacterial inhibitors from the sponge-derived fungus *Aspergillus* sp.. *Tetrahedron Letters*, 2014, 55(17): 2789-2792.
- [37] Song F, Ren B, Chen C, Yu K, Liu X, Zhang Y, Yang N, He H, Liu X, Dai H, Zhang L. Three new sterigmatocystin analogues from marine-derived fungus *Aspergillus versicolor* MF359. *Applied Microbiology and Biotechnology*, 2014, 98(8): 3753-3758.
- [38] Hiort J, Maksimenka K, Reichert M, Perovic-Ottstadt S, Lin WH, Wray V, Steube K, Schaumann K, Weber H, Proksch P. New natural products from the sponge-derived fungus *Aspergillus niger*. *Journal of Natural Products*, 2004, 67(9): 1532-1543.
- [39] Ui H, Shiomi K, Yamaguchi Y, Masuma R, Nagamitsu T, Takano D, Sunazuka T, Namikoshi M, Omura S. Nafuredin, a novel inhibitor of NADH-fumarate reductase, produced by *Aspergillus niger* FT-0554. *The Journal of Antibiotics (Tokyo)*, 2001, 54(3): 234-238.
- [40] Parvatkar RR, D' Souza, Tripathi A. Aspernolides A and B,

- butenolides from a marine-derived fungus *Aspergillus terreus*. *Phytochemistry*, 2009, 70(1): 128-132.
- [41] Wei MY, Wang CY, Liu QA, Shao CL, She ZG, Lin YC. Five sesquiterpenoids from a marine-derived fungus *Aspergillus* sp. isolated from a gorgonian *Dichotella gemmacea*. *Marine Drugs*, 2010, 8(4): 941-949.
- [42] Shao CL, Wang CY, Wei MY, Gu YC, She ZG, Qian PY, Lin YC. Aspergilones A and B, two benzylazaphilones with an unprecedented carbon skeleton from the gorgonian-derived fungus *Aspergillus* sp.. *Bioorganic and Medicinal Chemistry Letters*, 2011, 21(2): 690-693.
- [43] Trisuwan K, Rukachaisirikul V, Kaewpet M, Phongpaichit S, Hutadilok-Towatana N, Preedanon S, Sakayaroj J. Sesquiterpene and xanthone derivatives from the sea fan-derived fungus *Aspergillus sydowii* PSU-F154. *Journal of Natural Products*, 2011, 74(7): 1663-1667.
- [44] Zhuang Y, Teng X, Wang Y, Liu P, Wang H, Li J, Li G, Zhu W. Cyclopeptides and polyketides from coral-associated fungus, *Aspergillus versicolor* LCJ-5-4. *Tetrahedron*, 2011, 67(37): 7085-7089.
- [45] Zhuang Y, Teng X, Wang Y, Liu P, Li G, Zhu W. New quinazolinone alkaloids within rare amino acid residue from coral-associated fungus, *Aspergillus versicolor* LCJ-5-4. *Organic Letters*, 2011, 13(5): 1130-1133.
- [46] Hill RA, Sutherland A. Hot off the press. *Natural Products Report*, 2011, 28(10): 1621.
- [47] Hill RA, Sutherland A. Hot off the press. *Natural Products Report*, 2011, 28(6): 1031.
- [48] Wang L, Zhu W. Versicolactones A and B: total synthesis and structure revision. *Tetrahedron Letters*, 2013, 54(49): 6729-6731.
- [49] Zhang D, Satake M, Fukuzawa S, Sugahara K, Niitsu A, Shirai T, Tachibana K. Two new indole alkaloids, 2-(3, 3-dimethylprop-1-ene)-costaclavine and 2-(3, 3-dimethylprop-1-ene)-epicostaclavine, from the marine-derived fungus *Aspergillus fumigatus*. *Natural Product Communications*, 2012, 66(1): 222-226.
- [50] Afiyatullof SS, Zhuravleva OI, Chaikina EL, Anisimov MM. A new spirotryprostatin from the marine isolate of the fungus *Aspergillus fumigatus*. *Chemistry of Natural Compounds*, 2012, 48(1): 95-98.
- [51] Afiyatullof S, Zhuravleva OI, Antonov AS, Kalinovsky AI, Pivkin MV, Menchinskaya ES, Aminin DL. New metabolites from the marine-derived fungus *Aspergillus fumigatus*. *Natural Product Communications*, 2012, 7(4): 497-500.
- [52] He F, Sun YL, Liu KS, Zhang XY, Qian PY, Wang YF, Qi SH. Indole alkaloids from marine-derived fungus *Aspergillus sydowii* SCSIO 00305. *The Journal of Antibiotics (Tokyo)*, 2012, 65(2): 109-111.
- [53] Zhang D, Fukuzawa S, Satake M, Li X, Kuranaga T, Niitsu A, Yoshizawa K, Tachibana K. Ophiobolin O and 6-epi-ophiobolin O, two new cytotoxic sesterterpenes from the marine derived fungus *Aspergillus* sp.. *Natural Product Communications*, 2012, 7(11): 1411-1414.
- [54] Yang T, Lu Z, Meng L, Wei S, Hong K, Zhu W, Huang C. The novel agent ophiobolin O induces apoptosis and cell cycle arrest of MCF-7 cells through activation of MAPK signaling pathways. *Bioorganic and Medicinal Chemistry Letters*, 2012, 22(1): 579-585.
- [55] He F, Bao J, Zhang XY, Tu ZC, Shi YM, Qi SH. Asperterrestide A, a cytotoxic cyclic tetrapeptide from the marine-derived fungus *Aspergillus terreus* SCSGAF0162. *Journal of Natural Products*, 2013, 76(6): 1182-1186.
- [56] Chen M, Shao CL, Fu XM, Xu RF, Zheng JJ, Zhao DL, She ZG, Wang CY. Bioactive indole alkaloids and phenyl ether derivatives from a marine-derived *Aspergillus* sp. fungus. *Journal of Natural Products*, 2013, 76(4): 547-553.
- [57] Zheng CJ, Shao CL, Wu LY, Chen M, Wang KL, Zhao DL, Sun XP, Chen GY, Wang CY. Bioactive phenylalanine derivatives and cytochalasins from the soft coral-derived fungus, *Aspergillus elegans*. *Marine Drugs*, 2013, 11(6): 2054-2068.
- [58] Bao J, Xu XY, Zhang XY, Qi SH. A new macrolide from a marine-derived fungus *Aspergillus* sp.. *Natural Product Communications*, 2013, 8(8): 1127-1128.
- [59] Liu Z, Xia G, Chen S, Liu Y, Li H, She Z. Eurothiocin A and B, sulfur-containing benzofurans from a soft coral-derived fungus *Eurotium rubrum* SH-823. *Marine Drugs*, 2014, 12(6): 3669-3680.
- [60] Chen M, Fu X, Kong C, Wang C. Nucleoside derivatives from the marine-derived fungus *Aspergillus versicolor*. *Natural Product Research*, 2014, 28(12): 895-900.
- [61] Jiang T, Li T, Li J, Fu H, Pei Y, Lin W. Cerebroside analogues from marine-derived fungus *Aspergillus flavipes*. *Journal of Asian Natural Products Research*, 2004, 6(4): 249-257.
- [62] Snider BB, Zeng H. Total syntheses of (-)-fumiquinolines

- A, B, and I. *Organic Letters*, 2000, 2(25): 4103-4106.
- [63] Wang H, Ganesan A. Total synthesis of the fumiquinazoline alkaloids: solution-phase studies I. *Journal of Organic Chemistry*, 2000, 65(4): 1022-1030.
- [64] Takahashi C, Matsushita T, Doi M, Minoura K, Shingu T, Kumeda Y, Numata A. Fumiquinazolines A- G, novel metabolites of a fungus separated from a *Pseudolabrus* marine fish. *Journal of the Chemical Society, Dalton Transactions* 1, 1995, (18): 2345-2353.
- [65] Yamada T, Imai E, Nakatsuji K, Numata A, Tanaka R. Cephalimysin A, a potent cytotoxic metabolite from an *Aspergillus* species separated from a marine fish. *Tetrahedron Letters*, 2007, 48(36): 6294-6296.
- [66] Yamada T, Kitada H, Kajimoto T, Numata A, Tanaka R. The relationship between the CD cotton effect and the absolute configuration of FD-838 and its seven stereoisomers. *Journal of Organic Chemistry*, 2010, 75(12): 4146-4153.
- [67] Zhu T, Chen Z, Liu P, Wang Y, Xin Z, Zhu W. New rubrolides from the marine-derived fungus *Aspergillus terreus* OUCMDZ-1925. *The Journal of Antibiotics (Tokyo)*, 2014, 67(4): 315-318.
- [68] Tsukamoto S, Hirota H, Imachi M, Fujimuro M, Onuki H, Ohta T, Yokosawa H. Himeic acid A: a new ubiquitin-activating enzyme inhibitor isolated from a marine-derived fungus, *Aspergillus* sp.. *Bioorganic and Medicinal Chemistry Letters*, 2005, 15(1): 191-194.
- [69] Kuwana T, Miyazaki M, Kato H, Tsukamoto S. Himeic acids E- G, new 4-pyridone derivatives from a culture of *Aspergillus* sp.. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2013, 61(1): 105-107.
- [70] Kato H, Yoshida T, Tokue T, Nojiri Y, Hirota H, Ohta T, Williams RM, Tsukamoto S. Notoamides A- D: prenylated indole alkaloids isolated from a marine-derived fungus, *Aspergillus* sp.. *Angewandte Chemie International Edition*, 2007, 46(13): 2254-2256.
- [71] Tsukamoto S, Kato H, Samizo M, Nojiri Y, Onuki H, Hirota H, Ohta T. Notoamides F-K, prenylated indole alkaloids isolated from a marine-derived *Aspergillus* sp.. *Journal of Natural Products*, 2008, 71(12): 2064-2067.
- [72] Tsukamoto S, Kato H, Greshock TJ, Hirota H, Ohta T, Williams RM. Isolation of notoamide E, a key precursor in the biosynthesis of prenylated indole alkaloids in a marine-derived fungus, *Aspergillus* sp.. *Journal of the American Chemical Society*, 2009, 131(11): 3834-3835.
- [73] Tsukamoto S, Kawabata T, Kato H, Greshock TJ, Hirota H, Ohta T, Williams RM. Isolation of antipodal (-)-versicolamide B and notoamides L-N from a marine-derived *Aspergillus* sp.. *Organic Letters*, 2009, 11(6): 1297-1300.
- [74] Tsukamoto S, Umaoka H, Yoshikawa K, Ikeda T, Hirota H. Notoamide O, a structurally unprecedented prenylated indole alkaloid, and notoamides P-R from a marine-derived fungus, *Aspergillus* sp.. *Journal of Natural Products*, 2010, 73(8): 1438-1440.
- [75] Ding Y, de Wet JR, Cavalcoli J, Li S, Greshock TJ, Miller KA, Finefield JM, Sunderhaus JD, McAfoos TJ, Tsukamoto S, Williams RM, Sherman DH. Genome-based characterization of two prenylation steps in the assembly of the stephacidin and notoamide anticancer agents in a marine-derived *Aspergillus* sp.. *Journal of the American Chemical Society*, 2010, 132(36): 12733-12740.
- [76] Kitano M, Yamada T, Amagata T, Minoura K, Tanaka R, Numata A. Novel pyridino- α -pyrone sesquiterpene type pileotin produced by a sea urchin-derived *Aspergillus* sp.. *Tetrahedron Letters*, 2012, 53(32): 4192-4194.
- [77] Nakanishi K, Doi M, Usami Y, Amagata T, Minoura K, Tanaka R, Numata A, Yamada T. Anthcolorins A-F, novel cytotoxic metabolites from a sea urchin-derived *Aspergillus versicolor*. *Tetrahedron*, 2013, 69(23): 4617-4623.
- [78] Wang R, Liu T, Shen M, Yang M, Feng Q, Tang X, Li X. Spiculisporic acids B- D, three new γ -butenolide derivatives from a sea urchin-derived fungus *Aspergillus* sp. HDf2. *Molecules*, 2012, 17(11): 13175-13182.
- [79] Jiang W, Ye P, Chen CA, Wang K, Liu P, He S, Wu X, Gan L, Ye Y, Wu B. Two novel hepatocellular carcinoma cycle inhibitory cyclodepsipeptides from a hydrothermal vent crab-associated fungus *Aspergillus clavatus* C2WU. *Marine Drugs*, 2013, 11(12): 4761-4772.
- [80] Ye P, Shen L, Jiang W, Ye Y, Chen CA, Wu X, Wang K, Wu B. Zn-driven discovery of a hydrothermal vent fungal metabolite clavatuside C, and an experimental study of the anti-cancer mechanism of clavatuside B. *Marine Drugs*, 2014, 12(6): 3203-3217.
- [81] Wang F, Fang Y, Zhu T, Zhang M, Lin A, Gu Q, Zhu W. Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*. *Tetrahedron*, 2008, 64(34): 7986-7991.

- [82] Wang F, Li D, Zhu T, Zhang M, Gu Q. Pseurotin A1 and A2, two new 1-oxa-7-azaspiro [4. 4] non-2-ene-4, 6-diones from the holothurian-derived fungus *Aspergillus fumigatus* WFZ-25. *Canadian Journal of Chemistry*, 2011, 89(1): 72-76.
- [83] Bugni TS, Abbanat D, Bernan VS, Maiese WM, Greenstein M, Van Wagoner RM, Ireland CM. Yanuthones: novel metabolites from a marine isolate of *Aspergillus niger*. *Journal of Organic Chemistry*, 2000, 65(21): 7195-7200.
- [84] Tsukamoto S, Miura S, Yamashita Y, Ohta T. Aspermytin A: a new neurotrophic polyketide isolated from a marine-derived fungus of the genus *Aspergillus*. *Bioorganic and Medicinal Chemistry Letters*, 2004, 14(2): 417-420.
- [85] Sun K, Li Y, Guo L, Wang Y, Liu P, Zhu W. Indole diterpenoids and isocoumarin from the fungus, *Aspergillus flavus*, isolated from the Prawn, *Penaeus vannamei*. *Marine Drugs*, 2014, 12(7): 3970-3981.
- [86] Xu J, Zhao S, Yang X. A new cyclopeptide metabolite of marine gut fungus from *Ligia oceanica*. *Natural Product Research*, 2014, 28(13): 994-997.
- [87] Xu J, Nakazawa T, Ukai K, Kobayashi H, Mangindaan REP, Wewengkang DS, Rotinsulu H, Namikoshi M. Tetrahydrobostrycin and 1-deoxytetrahydrobostrycin, two new hexahydroanthrone derivatives, from a marine-derived fungus *Aspergillus* sp.. *The Journal of Antibiotics (Tokyo)*, 2008, 61(7): 415-419.
- [88] Rahbæk L, Christophersen C, Frisvad J, Bengaard HS, Larsen S, Rassing BR. Insulicolide A: a new nitrobenzoyloxy-substituted sesquiterpene from the marine fungus *Aspergillus insulicola*. *Journal of Natural Products*, 1997, 60(8): 811-813.
- [89] Belofsky GN, Jensen PR, Renner MK, Fenical W. New cytotoxic sesquiterpenoid nitrobenzoyl esters from a marine isolate of the fungus *Aspergillus versicolor*. *Tetrahedron*, 1998, 54(9): 1715-1724.
- [90] Lorenz P, Jensen PR, Fenical W. Mactanamide, a new fungistatic diketopiperazine produced by a marine *Aspergillus* sp.. *Natural Products Letters*, 1998, 12(1): 55-60.
- [91] Son BW, Choi JS, Kim JC, Nam KW, Kim D, Chung HY, Kang JS, Choi HD. Parasitenone, a new epoxycyclohexenone related to gabosine from the marine-derived fungus *Aspergillus parasiticus*. *Journal of Natural Products*, 2002, 65(5): 794-795.
- [92] Lee SM, Li XF, Jiang H, Cheng JG, Seong S, Choi HD, Son BW. Terreusinone, a novel UV-A protecting dipyrroloquinone from the marine algicolous fungus *Aspergillus terreus*. *Tetrahedron Letters*, 2003, 44(42): 7707-7710.
- [93] Li Y, Li X, Kim S, Kang JS, Choi HD, Rho JR, Son BW. Golmaenone, a new diketopiperazine alkaloid from the marine-derived fungus *Aspergillus* sp.. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2004, 52(3): 375-376.
- [94] Teuscher F, Lin W, Wray V, Edrada R, Padmakumar K, Proksch P, Ebel R. Two new cyclopentanoids from the endophytic fungus *Aspergillus sydowii* associated with the marine alga *Acanthophora spicifera*. *Natural Product Communications*, 2006, 1(11): 927-933.
- [95] Nguyen HP, Zhang D, Lee U, Kang JS, Choi HD, Son BW. Dehydroxychlorofusarielin B, an antibacterial polyoxygenated decalin derivative from the marine-derived fungus *Aspergillus* sp.. *Journal of Natural Products*, 2007, 70(7): 1188-1190.
- [96] Zhang Y, Li X, Wang B. Nigerasperones A- C, new monomeric and dimeric naphtho- γ -pyrones from a marine alga-derived endophytic fungus *Aspergillus niger* EN-13. *The Journal of Antibiotics (Tokyo)*, 2007, 60(3): 204-210.
- [97] Zhang Y, Wang S, Li XM, Cui CM, Feng C, Wang BG. New sphingolipids with a previously unreported 9-methyl-C20-sphingosine moiety from a marine alga-derived endophytic fungus *Aspergillus niger* EN-13. *Lipids*, 2007, 42(8): 759-764.
- [98] Zhang Y, Li X, Proksch P, Wang B. Ergosterimide, a new natural diels- alder adduct of a steroid and maleimide in the fungus *Aspergillus niger*. *Steroids*, 2007, 72(9): 723-727.
- [99] Zhang Y, Li XM, Wang CY. A new naphthoquinoneimine derivative from the marine alga-derived endophytic fungus *Aspergillus niger* EN-13. *Chinese Chemical Letters*, 2007, (8): 951-953.
- [100] Zhang Y, Li XM, Feng Y, Wang BG. Phenethyl- α -pyrone derivatives and cyclodipeptides from a marine alga-derived endophytic fungus *Aspergillus niger* EN-13. *Natural Product Research*, 2010, 24(11): 1036-1043.
- [101] Lin A, Lu X, Fang Y, Zhu T, Gu Q, Zhu W. Two new 5-hydroxy-2-pyrone derivatives isolated from a marine-derived fungus *Aspergillus flavus*. *The Journal of Antibiotics (Tokyo)*, 2008, 61(4): 245-249.
- [102] Lin A, Du L, Fang Y, Wang F, Zhu T, Gu Q, Zhu W. Iso- α -cyclopiazonic acid, a new natural product isolated from the marine-derived fungus *Aspergillus flavus* C-F-3. *Chemistry of Natural Compounds*, 2009, (5): 677-680.
- [103] Cui CM, Li XM, Li CS, Sun HF, Gao SS, Wang BG.

- Benzodiazepine alkaloids from marine-derived endophytic fungus *Aspergillus ochraceus*. *Helvetica Chimica Acta*, 2009, 92(7): 1366-1370.
- [104] Cui CM, Li XM, Meng L, Li CS, Huang CG, Wang BG. 7-Nor-ergosterolide, a pentalactone-containing norsteroid and related steroids from the marine-derived endophytic *Aspergillus ochraceus* EN-31. *Journal of Natural Products*, 2010, 73(11): 1780-1784.
- [105] Qiao MF, Ji NY, Liu XH, Li K, Zhu QM, Xue QZ. Indoloditerpenes from an algicolous isolate of *Aspergillus oryzae*. *Bioorganic and Medicinal Chemistry Letters*, 2010, 20(19): 5677-5680.
- [106] Qiao MF, Ji NY, Liu XH, Li F, Xue QZ. Asporierygosterol, a new steroid from an algicolous isolate of *Aspergillus oryzae*. *Natural Product Communications*, 2010, 5(10): 1575-1578.
- [107] Izumikawa M, Hashimoto J, Takagi M, Shin-ya K. Isolation of two new terpeptin analogs—JBIR-81 and JBIR-82—from a seaweed-derived fungus, *Aspergillus* sp. SpD081030G1f1. *The Journal of Antibiotics (Tokyo)*, 2010, 63(7): 389-391.
- [108] Yang G, Sandjo L, Yun K, Leutou AS, Kim GD, Choi HD, Kang JS, Hong J, Son BW. Flavusides A and B, antibacterial cerebroside from the marine-derived fungus *Aspergillus flavus*. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2011, (9): 1174-1177.
- [109] Qiao MF, Ji NY, Miao FP, Yin XL. Steroids and an oxylipin from an algicolous isolate of *Aspergillus flavus*. *Magnetic Resonance in Chemistry*, 2011, 49(6): 366-369.
- [110] Zhuravleva OI, Afyatullov S, Denisenko VA, Ermakova SP, Slinkina NN, Dmitrenok PS, Kim NY. Secondary metabolites from a marine-derived fungus *Aspergillus carneus* Blochwitz. *Phytochemistry*, 2012, 80: 123-131.
- [111] Zhuravleva OI, Afyatullov S, Yurchenko EA, Denisenko VA, Kirichuk NN, Dmitrenok PS. New metabolites from the algal associated marine-derived fungus *Aspergillus carneus*. *Natural Product Communications*, 2013, 8(8): 1071-1074.
- [112] Sun HF, Li XM, Meng L, Cui CM, Gao SS, Li CS, Huang CG, Wang BG. Asperolides A-C, tetranorlabdane diterpenoids from the marine alga-derived endophytic fungus *Aspergillus wentii* EN-48. *Journal of Natural Products*, 2012, 75(2): 148-152.
- [113] Li X, Li X, Xu G, Li C, Wang B. Antioxidant metabolites from marine alga-derived fungus *Aspergillus wentii* EN-48. *Phytochemistry Letters*, 2014, 7: 120-123.
- [114] Hawas UW, El-Beih AA, El-Halawany AM. Bioactive anthraquinones from endophytic fungus *Aspergillus versicolor* isolated from red sea algae. *Archives of Pharmacal Research*, 2012, 35(10): 1749-1756.
- [115] Liu XH, Miao FP, Qiao MF, Cichewicz RH, Ji NY. Terretinin, ophiobolin, and drimane terpenes with absolute configurations from an algicolous *Aspergillus ustus*. *RSC Advances*, 2013, 3(2): 588-595.
- [116] Miao F, Li X, Liu X, Cichewicz RH, Ji N. Secondary metabolites from an algicolous *Aspergillus versicolor* strain. *Marine Drugs*, 2012, 10(1): 131-139.
- [117] Liu X, Miao F, Li X, Yin X, Ji N. A new sesquiterpene from an endophytic *Aspergillus versicolor* strain. *Natural Product Communications*, 2012, 7(7): 819-820.
- [118] Ji N, Liu X, Miao F, Qiao M. Aspeverin, a new alkaloid from an algicolous strain of *Aspergillus versicolor*. *Organic Letters*, 2013, 15(10): 2327-2329.
- [119] Zhang Y, Li XM, Wang BG. Anthraquinone derivatives produced by marine-derived fungus *Aspergillus versicolor* EN-7. *Bioscience, Biotechnology and Biochemistry*, 2012, (9): 1774-1776.
- [120] Li Y, Sun K, Wang Y, Fu P, Liu P, Wang C, Zhu W. A cytotoxic pyrrolidinoindoline diketopiperazine dimer from the algal fungus *Eurotium herbariorum* HT-2. *Chinese Chemical Letters*, 2013, 24(12): 1049-1052.
- [121] Fang W, Lin X, Zhou X, Wan J, Lu X, Yang B, Ai W, Lin J, Zhang T, Tu Z. Cytotoxic and antiviral nitrobenzoyl sesquiterpenoids from the marine-derived fungus *Aspergillus ochraceus* Jcma1F17. *Medicinal Chemistry Communications*, 2014, 5(6): 701-705.
- [122] Ebada SS, Fischer T, Hamacher A, Du F, Roth YO, Kassack MU, Wang B, Roth EH. Psychrophilin E, a new cyclotriptide, from co-fermentation of two marine alga-derived fungi of the genus *Aspergillus*. *Natural Product Research*, 2014, 28(11): 776-781.
- [123] Ogawa A, Murakami C, Kamisuki S, Kuriyama I, Yoshida H, Sugawara F, Mizushima Y. Pseudodelectusin, a novel isochroman derivative from *Aspergillus pseudodelectus* a parasite of the sea weed, *Sargassum fusiform*, as a selective human cancer cytotoxin. *Bioorganic and Medicinal Chemistry Letters*, 2004, 14(13): 3539-3543.
- [124] Kanoh K, Kohno S, Asari T, Harada T, Katada J, Muramatsu M, Kawashima H, Sekiya H, Uno I. (-)-Phenylahistin: a new

- mammalian cell cycle inhibitor produced by *Aspergillus ustus*. *Bioorganic and Medicinal Chemistry Letters*, 1997, 7(22): 2847-2852.
- [125] Couladouros EA, Magos AD. Solid-phase total synthesis of (–)-phenylhistine and (–)-aurantiamine. Synthesis of a diverse dehydro-2, 5-diketopiperazine library. Part II. *Molecular Diversity*, 2005, 9(1-3): 111-121.
- [126] Shangguan N, Joullié MM. Total synthesis of isoroquefortine E and phenylahistin. *Tetrahedron Letters*, 2009, 50(49): 6755-6757.
- [127] Yakushiji F, Tanaka H, Muguruma K, Iwahashi T, Yamazaki Y, Hayashi Y. Prodrug study of plinabulin using a click strategy focused on the effects of a replaceable water-solubilizing moiety. *Chemical and Pharmaceutical Bulletin*, 2012, 60(7): 877-881.
- [128] Mita MM, Spear MA, Yee LK, Mita AC, Heath EI, Papadopoulos KP, Federico KC, Reich SD, Romero O, Malburg L. Phase 1 first-in-human trial of the vascular disrupting agent plinabulin (NPI-2358) in patients with solid tumors or lymphomas. *Clinical Cancer Research*, 2010, 16(23): 5892-5899.
- [129] Millward M, Mainwaring P, Mita A, Federico K, Lloyd GK, Reddinger N, Nawrocki S, Mita M, Spear MA. Phase 1 study of the novel vascular disrupting agent plinabulin (NPI-2358) and docetaxel. *Investigational New Drugs*, 2012, 30(3): 1065-1073.
- [130] Hayashi Y, Takeno H, Chinen T, Muguruma K, Okuyama K, Taguchi A, Takayama K, Yakushiji F, Miura M, Usui T. Development of a new benzophenone-diketopiperazine-type potent anti-microtubule agent possessing a 2-pyridine structure. *ACS Medicinal Chemistry Letters*, 2014, 5(10): 1094-1098.
- [131] Schlingmann G, Taniguchi T, He H, Bigelis R, Yang HY, Koehn FE, Carter GT, Berova N. Reassessing the structure of pyranonigrin. *Journal of Natural Products*, 2007, 70(7): 1180-1187.
- [132] Lin Z, Zhang G, Zhu T, Liu R, Wei H, Gu Q. Bioactive cytochalasins from *Aspergillus flavipes*, an endophytic fungus associated with the mangrove plant *Acanthus ilicifolius*. *Helvetica Chimica Acta*, 2009, 92(8): 1538-1544.
- [133] Li D, Li X, Wang B. Natural anthraquinone derivatives from a marine mangrove plant-derived endophytic fungus *Eurotium rubrum*: structural elucidation and DPPH radical scavenging activity. *Journal of Industrial Microbiology and Biotechnology*, 2009, 19(7): 675-680.
- [134] Huang HB, Feng XJ, Liu L, Chen B, Lu YJ, Ma L, She ZG, Lin YC. Three dimeric naphtho-gamma-pyrones from the mangrove endophytic fungus *Aspergillus tubingensis* isolated from *Pongamia pinnata*. *Planta Medica*, 2010, 76(16): 1888-1891.
- [135] Huang HB, Xiao ZE, Feng XJ, Huang CH, Zhu X, Ju JH, Li MF, Lin YC, Liu L, She ZG. Cytotoxic naphtho-g-pyrones from the mangrove endophytic fungus *Aspergillus tubingensis* (GX1-5E). *Helvetica Chimica Acta*, 2011, 94(9): 1732-1740.
- [136] Liu D, Li XM, Meng L, Li CS, Gao SS, Shang Z, Proksch P, Huang CG, Wang BG. Nigerapyrones A-H, alpha-pyrone derivatives from the marine mangrove-derived endophytic fungus *Aspergillus niger* MA-132. *Journal of Natural Products*, 2011, 74(8): 1787-1791.
- [137] Song YX, Qiao LT, Wang JJ, Zeng HM, She ZG, Miao CD, Hong K, Gu YC, Liu L, Lin YC. Two new meroterpenes from the mangrove endophytic fungus *Aspergillus* sp. 085241B. *Helvetica Chimica Acta*, 2011, 94(10): 1875-1880.
- [138] Wang H, Lu Z, Qu H, Liu P, Miao C, Zhu T, Li J, Hong K, Zhu W. Antimicrobial aflatoxins from the marine-derived fungus *Aspergillus flavus* 092008. *Archives of Pharmacol Research*, 2012, 35(8): 1387-1392.
- [139] Li S, Wei M, Chen G, Lin Y. Two new dihydroisocoumarins from the endophytic fungus *Aspergillus* sp. collected from the south china sea. *Chemistry of Natural Compounds*, 2012, 48(3): 371-373.
- [140] An C, Li X, Luo H, Li C, Wang M, Xu G, Wang B. 4-phenyl-3, 4-dihydroquinolone derivatives from *Aspergillus nidulans* MA-143, an endophytic fungus isolated from the mangrove plant *Rhizophora stylosa*. *Journal of Natural Products*, 2013, 76(10): 1896-1901.
- [141] An C, Li X, Li C, Wang M, Xu G, Wang B. Aniquinazolines A- D, four new quinazolinone alkaloids from marine-derived endophytic fungus *Aspergillus nidulans*. *Marine Drugs*, 2013, 11(7): 2682-2694.
- [142] Bai Z, Lin X, Wang Y, Wang J, Zhou X, Yang B, Liu J, Yang X, Wang Y, Liu Y. New phenyl derivatives from endophytic fungus *Aspergillus flavipes* AIL8 derived of mangrove plant *Acanthus ilicifolius*. *Fitoterapia*, 2014, 95: 194-202.
- [143] Xiao Z, Huang H, Shao C, Xia X, Ma L, Huang X, Lu Y, Lin Y, Long Y, She Z. Asperterpenols A and B, new

- sesterterpenoids isolated from a mangrove endophytic fungus *Aspergillus* sp. 085242. *Organic Letters*, 2013, 15(10): 2522-2525.
- [144] Huang X, Huang H, Li H, Sun X, Huang H, Lu Y, Lin Y, Long Y, She Z. Asperterpenoid A, a new sesterterpenoid as an inhibitor of mycobacterium tuberculosis protein tyrosine phosphatase B from the culture of *Aspergillus* sp. 16-5c. *Organic Letters*, 2013, 15(4): 721-723.
- [145] Deng C, Liu S, Huang C, Pang J, Lin Y. Secondary metabolites of a mangrove endophytic fungus *Aspergillus terreus* (No. GX7-3B) from the South China Sea. *Marine Drugs*, 2013, 11(7): 2616-2624.
- [146] Tsuda M, Mugishima T, Komatsu K, Sone T, Tanaka M, Mikami Y, Shiro M, Hirai M, Ohizumi Y, Kobayashi J. Speradine A, a new pentacyclic oxindole alkaloid from a marine-derived fungus *Aspergillus tamarii*. *Tetrahedron*, 2003, 59(18): 3227-3230.
- [147] Zhang M, Wang W, Fang Y, Zhu T, Gu Q, Zhu W. Cytotoxic alkaloids and antibiotic nordammarane triterpenoids from the marine-derived fungus *Aspergillus sydowi*. *Journal of Natural Products*, 2008, 71(6): 985-989.
- [148] Capon RJ, Skene C, Stewart M, Ford J, Richard AJ, Williams L, Lacey E, Gill JH, Heiland K, Friedel T. Aspergillicins A-E: five novel depsipeptides from the marine-derived fungus *Aspergillus carneus*. *Organic and Biomolecular Chemistry*, 2003, 1(11): 1856-1862.
- [149] Smetanina OF, Kalinovskii AI, Khudyakov YV, Moiseenko OP, Pivkin MV, Menzorova NI, Sibirtsev YT, Kuznetsova TA. Metabolites of the marine fungus *Aspergillus varians* KMM 4630. *Chemistry of Natural Compounds*, 2005, 41(2): 243-244.
- [150] Liu R, Zhu W, Zhang Y, Zhu T, Liu H, Fang Y, Gu Q. A new diphenyl ether from marine-derived fungus *Aspergillus* sp. BF-2. *The Journal of Antibiotics (Tokyo)*, 2006, 59(6): 362-365.
- [151] Boot CM, Gassner NC, Compton JE, Tenney K, Tamble CM, Lokey RS. Pinpointing pseurotins from a marine-derived *Aspergillus* as tools for chemical genetics using a synthetic lethality yeast screen. *Journal of Natural Products*, 2007, 70(10): 1672-1675.
- [152] Kamigauchi T, Sakazaki R, Nagashima K, Kawamura Y, Yasuda Y, Matsushima K, Tani H, Takahashi Y, Ishii K, Suzuki R, Koizumi K, Nakai H, Ikenishi Y, Terui Y. Terpenins, novel immunosuppressants produced by *Aspergillus candidus*. *The Journal of Antibiotics (Tokyo)*, 1998, 51(4): 445-450.
- [153] Zhang Y, Zhu T, Fang Y, Liu H, Gu Q, Zhu W. Carbonarones A and B, new bioactive γ -pyrone and α -pyridone derivatives from the marine-derived fungus *Aspergillus carbonarius*. *The Journal of Antibiotics (Tokyo)*, 2007, 60(2): 153-157.
- [154] Zhang Y, Ling S, Fang Y, Zhu T, Gu Q, Zhu W. Isolation, structure elucidation and antimycobacterial properties of dimeric naphtho- γ -pyrones from the marinederived fungus *Aspergillus carbonarius*. *Chemistry & Biodiversity*, 2008, 5(1): 93-100.
- [155] Zhao WY, Zhu TJ, Han XX, Fan GT, Liu HB. A new gliotoxin analogue from a marine-derived fungus *Aspergillus fumigatus* Fres. *Natural Product Research*, 2009, 23(3): 203-207.
- [156] Zhao WY, Zhu TJ, Fan GT, Liu HB, Fang YC. Three new dioxopiperazine metabolites from a marine-derived fungus *Aspergillus fumigatus* Fres. *Natural Product Research*, 2010, 24(10): 953-957.
- [157] Fremlin LJ, Piggott AM, Lacey E, Capon RJ. Cottoquinazoline A and cotteslosins A and B, metabolites from an Australian marine-derived strain of *Aspergillus versicolor*. *Journal of Natural Products*, 2009, 72(4): 666-670.
- [158] Ren H, Liu R' CL, Zhu T, Zhu W, Gu Q. Two new hetero-spirocyclic gamma-lactam derivatives from marine sediment-derived fungus *Aspergillus sydowi* D2-6. *Archives of Pharmacol Research*, 2010, 33(4): 499-502.
- [159] Crews MS, Draskovic M, Sohn J, Johnson TA, Tenney K, Valeriote FA, Yao X. Azonazine, a novel dipeptide from a Hawaiian marine sediment-derived fungus, *Aspergillus insulicola*. *Organic Letters*, 2010, 12(20): 4458-4461.
- [160] Yurchenko AN, Smetanina OF, Kalinovskiy AI, Pivkin MV, Dmitrenok PS, Kuznetsova TA. A new meroterpenoid from the marine fungus *Aspergillus versicolor* (Vuill.) Tirab. *Russian Chemical Bulletin*, 2010, 59(4): 852-856.
- [161] Liu S, Shen Y. A new cyclic peptide from the marine fungal strain *Aspergillus* sp. AF119. *Chemistry of Natural Compounds*, 2011, 47(5): 786-788.
- [162] Choi EJ, Park JS, Kim YJ, Jung JH, Lee JK, Kwon HC, Yang HO. Apoptosisinducing effect of diketopiperazine disulfides produced by *Aspergillus* sp. KMD 901 isolated from marine sediment on HCT116 colon cancer cell lines. *Journal of*

- Applied Microbiology*, 2011, 110(1): 304-313.
- [163] Liu S, Lu C, Huang J, Shen Y. Three new compounds from the marine fungal strain *Aspergillus* sp. AF119. *Records of Natural Products*, 2012, 6(4): 334-338.
- [164] Liu S, Zhao B, Lu C, Huang J, Shen Y. Two new p-terphenyl derivatives from the marine fungal strain *Aspergillus* sp. AF119. *Natural Product Communications*, 2012, 7(8): 1057-1062.
- [165] Sohn J, Oh H. Protulactones A and B: Two new polyketides from the marine-derived fungus *Aspergillus* sp. SF-5044. *Bulletin of the Korean Chemical Society*, 2010, 31(6): 1695-1698.
- [166] Lee SU, Asami Y, Lee D, Jang JH, Ahn JS, Oh H. Protuboxepins A and B and protubonines A and B from the marine-derived fungus *Aspergillus* sp. SF-5044. *Journal of Natural Products*, 2011, 74(5): 1284-1287.
- [167] Neff SA, Lee SU, Asami Y, Ahn JS, Oh H, Baltrusaitis J, Gloer JB, Wicklow DT. Aflaquinolones A- G: secondary metabolites from marine and fungicolous isolates of *Aspergillus* sp.. *Journal of Natural Products*, 2012, 75(3): 464-472.
- [168] Zhou LN, Gao HQ, Cai SX, Zhu TT, Gu QQ, Li DH. Two new cyclic pentapeptides from the marine-derived fungus *Aspergillus versicolor*. *Helvetica Chimica Acta*, 2011, 94(6): 1065-1070.
- [169] Gao H, Zhou L, Cai S, Zhang G, Zhu T, Gu Q, Li D. Diorcinols BE, new prenylated diphenyl ethers from the marine-derived fungus *Aspergillus versicolor* ZLN-60. *The Journal of Antibiotics (Tokyo)*, 2013, 66(9): 539-542.
- [170] Cai S, Sun S, Zhou H, Kong X, Zhu T, Li D, Gu Q. Prenylated polyhydroxy-p-terphenyls from *Aspergillus taichungensis* ZHN-7-07. *Journal of Natural Products*, 2011, 74(5): 1106-1110.
- [171] Zhou H, Zhu T, Cai S, Gu Q, Li D. Drimane sesquiterpenoids from the mangrove-derived fungus *Aspergillus ustus*. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2011, 59(6): 762-766.
- [172] Wang Y. LZ, J Bai LMZ, X Wu LZ, Pei YH. 2,5-diketopiperazines from the marine-derived fungus *Aspergillus fumigatus* YK-7. *Chemistry and Biodiversity*, 2012, 9(2): 385-393.
- [173] Shen Y, Zou J, Xie D, Ge H, Cao X, Dai J. Butyrolactone and cycloheptanetrione from mangrove-associated fungus *Aspergillus terreus*. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2012, 60(11): 1437-1441.
- [174] Wu QX, Jin XJ, Draskovic M, Crews MS, Tenney K, Valeriote FA, Yao XJ, Crews P. Unraveling the numerous biosynthetic products of the marine sediment-derived fungus, *Aspergillus insulicola*. *Phytochemistry Letters*, 2012, 5(1): 114-117.
- [175] Zhuravleva OI, Afiyatulloev SS, Vishchuk OS, Denisenko VA, Slinkina NN, Smetanina OF. Decumbenone C, a new cytotoxic decaline derivative from the marine fungus *Aspergillus sulphureus* KMM 4640. *Archives of Pharmacal Research*, 2012, 35(10): 1757-1762.
- [176] Song F, Liu X, Guo H, Ren B, Chen C, Piggott AM, Yu K, Gao H, Wang Q, Liu M, Liu X, Dai H, Zhang L, Capon RJ. Brevianamides with antitubercular potential from a marine-derived isolate of *Aspergillus versicolor*. *Organic Letters*, 2012, 14(18): 4770-4773.
- [177] Lu Z, Wang Y, Miao C, Liu P, Hong K, Zhu W. Sesquiterpenoids and benzofuranoids from the marine-derived fungus *Aspergillus ustus* 094102. *Journal of Natural Products*, 2009, 72(10): 1761-1767.
- [178] Gao H, Liu W, Zhu T, Mo X, Má ndi A, Kurtá n T, Li J, Ai J, Gu Q, Li D. Diketopiperazine alkaloids from a mangrove rhizosphere soil derived fungus *Aspergillus effuses* H1-1. *Organic and Biomolecular Chemistry* 2012, 10(47): 9501-9506.
- [179] 刘为忠. 五株海洋真菌次生代谢产物及抗肿瘤活性研究. 中国海洋大学博士学位论文. 2006.
- [180] Peng J, Zhang X, Tu Z, Xu X, Qi S. Alkaloids from the deep-sea-derived fungus *Aspergillus westerdijkiae* DFFSCS013. *Journal of Natural Products*, 2013, 76(5): 983-987.
- [181] Liu X, Song F, Ma L, Chen C, Xiao X, Ren B, Liu X, Dai H, Piggott AM, Av-Gay Y. Sydowiols A- C: *Mycobacterium tuberculosis* protein tyrosine phosphatase inhibitors from an East China Sea marine-derived fungus, *Aspergillus sydowii*. *Tetrahedron Letters*, 2013, 54(45): 6081-6083.
- [182] Chen L, Zhang W, Zheng Q. Aculeatusquinones A-D, novel metabolites from the marine-derived fungus *Aspergillus aculeatus*. *Heterocycles*, 2013, 87(4): 861-868.
- [183] Peng JX, Gao HQ, Li J, Ai J, Geng MY, Zhang GJ, Zhu TT, Gu QQ, Li DH. Prenylated indole diketopiperazines from the marine- derived fungus *Aspergillus versicolor*. *The Journal of Organic Chemistry*, 2014, 79(17): 7895-7904.

- [184] Hu X, Xia Q, Zhao Y. Speradines B-E, four novel tetracyclic oxindole alkaloids from the marine-derived fungus *Aspergillus oryzae*. *Heterocycles*, 2014, 89(7): 1662-1669.
- [185] Hu X, Xia Q, Zhao Y, Zheng Q, Liu Q, Chen L, Zhang Q. Speradines F-H, three new oxindole alkaloids from the marine-derived fungus *Aspergillus Oryzae*. *Chemical and Pharmaceutical Bulletin (Tokyo)*, 2014, 62(9): 942-946.
- [186] Dai J, Carté BK, Sidebottom PJ, Sek Yew AL, Ng S, Huang Y, Butler MS. Circumdatin G, a new alkaloid from the fungus *Aspergillus ochraceus*. *Journal of Natural Products*, 2001, 64(1): 125-126.
- [187] Sun S, Ji C, Gu Q, Li D, Zhu T. Three new polyketides from marine-derived fungus *Aspergillus glaucus* HB1-19. *Journal of Asian Natural Product Research*, 2013, 15(9): 956-961.
- [188] Du L, Zhu TT, Fang YC, Liu HB, Gu QQ, Zhu WM. Aspergiolide A, a novel anthraquinone derivative with naphtho[1, 2, 3-de]chromene-2, 7-dione skeleton isolated from a marine-derived fungus *Aspergillus glaucus*. *Tetrahedron*, 2007, 63(5): 1085-1088.
- [189] Zheng J, Zhu H, Hong K, Wang Y, Liu P, Wang X, Peng X, Zhu W. Novel cyclic hexapeptides from marine-derived fungus, *Aspergillus sclerotiorum* PT06-1. *Organic Letters*, 2009, 11(22): 5262-5265.
- [190] Zheng J, Xu Z, Wang Y, Hong K, Liu P, Zhu W. Cyclic tripeptides from the halotolerant fungus *Aspergillus sclerotiorum* PT06-1. *Journal of Natural Products*, 2010, 73(6): 1133-1137.
- [191] Wang H, Zheng J, Qu H, Liu P, Wang Y, Zhu W. A new cytotoxic indole-3-ethenamide from the halotolerant fungus *Aspergillus sclerotiorum* PT06-1. *The Journal of Antibiotics (Tokyo)*, 2011, 64(10): 679-681.
- [192] Wang Y, Zheng J, Liu P, Wang W, Zhu W. Three new compounds from *Aspergillus terreus* PT06-2 grown in a high salt medium. *Marine Drugs*, 2011, 9(8): 1368-1378.
- [193] Cui C. Tryprostatins A and B, novel mammalian cell cycle inhibitors produced by *Aspergillus fumigatus*. *The Journal of Antibiotics (Tokyo)*, 1995, 48(11): 1382-1384.
- [194] Ouyang M, Liu R, Kuo Y. A new cerebroside, asperiamide A, from the marine fungus *Asperillus* sp. Note. *Journal of Asian Natural Product Research*, 2005, 7(5): 761-765.
- [195] Wu Z, Ouyang M, Su R, Guo Y. Two new cerebrosides and anthraquinone derivatives from the marine fungus *Aspergillus niger*. *Chinese Journal of Chemistry*, 2008, 26(4): 759-764.
- [196] Wu Z, Ouyang M, Tan Q. New asperxanthone and asperbiphenyl from the marine fungus *Aspergillus* sp.. *Pest Management Science*, 2009, 65(1): 60-65.
- [197] Dong Y, Cui C, Li C, Hua W, Wu C, Zhu T, Gu Q. Activation of dormant secondary metabolite production by introducing neomycin resistance into the deep-sea fungus, *Aspergillus versicolor* ZBY-3. *Marine Drugs*, 2014, 12(8): 4326-4352.
- [198] Cai S, Zhu T, Du L, Zhao B, Li D, Gu Q. Sterigmatocystins from the deep-sea-derived fungus *Aspergillus versicolor*. *The Journal of Antibiotics (Tokyo)*, 2010, 64(2): 193-196.
- [199] Kong X, Cai S, Zhu T, Gu Q, Li D, Luan Y. Secondary metabolites of a deep sea derived fungus *Aspergillus versicolor* CXCTD-06-6a and their bioactivity. *Journal of Ocean University of China*, 2014, 13(4): 691-695.
- [200] Cueto M, Jensen PR, Fenical W. Aspergilloxide, a novel sesterterpene epoxide from a marine-derived fungus of the genus *Aspergillus*. *Organic Letters*, 2002, 4(9): 1583-1585.
- [201] Ovenden SPB, Sberna G, Tait RM, Wildman HG, Patel R, Li B, Steffy K, Nguyen N, Meurer-Grimes BM. A diketopiperazine dimer from a marine-derived isolate of *Aspergillus niger*. *Journal of Natural Products*, 2004, 67(12): 2093-2095.
- [202] Yan WP, Song H, Song FH, Guo YS, Wu CH, Her, AS, Pu Y, Wang S, Naowarojna N, Weitz A, Hendrich MP, Costello CE, Zhang LX, Liu PH, Zhang YJ. Endoperoxide formation by an α -ketoglutarate-dependent mononuclear non-haem iron enzyme. *Nature*, 2015, 527: 539-543.
- [203] Xie X, Watanabe K, Wojcicki WA, Wang CC, Tang Y. Biosynthesis of lovastatin analogs with a broadly specific acyltransferase. *Chemistry & Biology*, 2006, 13(11): 1161-1169.
- [204] Mulder KC, Mulinari F, Franco OL, Soares MS, Magalhaes BS, Parachin NS. Lovastatin production: from molecular basis to industrial process optimization. *Biotechnology Advances*, 2015, 33(6 Pt 1): 648-665.
- [205] Xie X, Wong WW, Tang Y. Improving simvastatin bioconversion in *Escherichia coli* by deletion of *bioH*. *Metabolic Engineering*, 2007, 9(4): 379-386.
- [206] Xie X, Tang Y. Efficient synthesis of simvastatin using whole-cell biocatalysis. *Applied and Environmental Microbiology*, 2007, 73(7): 2054-2060.

New natural products from the marine-derived *Aspergillus* fungi-A review

Chengying Zhao, Haishan Liu, Weiming Zhu^{*}

Key Laboratory of Marine Drugs, Ministry of Education, School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, Shandong Province, China

Abstract: Marine-derived fungi were the main source of marine microbial natural products (NPs) due to their complex genetic background, chemodiversity and high yield of NPs. According to our previous survey for marine microbial NPs from 2010 to 2013, *Aspergillus* fungi have received the most of attention among all the marine-derived fungi, which accounted for 31% NPs of the marine fungal origins. This paper reviewed the sources, chemical structures and bioactivities of all the 512 new marine NPs of *Aspergillus* fungal origins from 1992 to 2014. These marine NPs have diverse chemical structures including polyketides, fatty acids, sterols and terpenoids, alkaloids, peptides, and so on, 36% of which displayed bioactivities such as cytotoxicity, antimicrobial activity, antioxidant and insecticidal activity. Nitrogen compounds are the major secondary metabolites accounting for 52% NPs from the marine-derived *Aspergillus* fungi. Nitrogen compounds are also the class with the highest ratio of bioactive compounds, 40% of which are bioactive. Plinabulin, a dehydrotetrapiperazine derivative of halimide had been ended its phase II trial and has received its phase III study from the third quarter of 2015 for the treatment of advanced, metastatic non-small cell lung cancer.

Keywords: marine-derived fungi, *Aspergillus* sp., natural products, chemical structures, bioactivities, sources

(本文责编: 张晓丽)

Supported by the National Natural Science Foundation of China (41376148, 81561148012, U1406402-1, U1501221)

*Corresponding author. Tel/Fax: +86-532-82031836; E-mail: weimingzhu@ouc.edu.cn

Received: 15 October 2015; Revised: 25 December 2015; Published Online: 30 December 2015